

NUMERICAL METHODS FOR FRACTIONAL DIFFUSION

ANDREA BONITO, JUAN PABLO BORTHAGARAY, RICARDO H. NOCHETTO,
ENRIQUE OTÁROLA, AND ABNER J. SALGADO

ABSTRACT. We present three schemes for the numerical approximation of fractional diffusion, which build on different definitions of such a non-local process. The first method is a PDE approach that applies to the spectral definition and exploits the extension to one higher dimension. The second method is the integral formulation and deals with singular non-integrable kernels. The third method is a discretization of the Dunford-Taylor formula. We discuss pros and cons of each method, error estimates, and document their performance with a few numerical experiments.

1. INTRODUCTION

Diffusion is the tendency of a substance to evenly spread into an available space, and is one of the most common physical processes. The classical models of diffusion, which usually start from the assumption of Brownian motion [1, 59], lead to well known models and even better studied equations. However, in recent times, it has become evident that many of the assumptions that lead to these models are not always satisfactory or even realistic at all. For this reason, new models of diffusion have been introduced. These, as a rule, are not based on the postulate that the underlying stochastic process is given by Brownian motion, so that the diffusion is regarded as anomalous [89]. The evidence of anomalous diffusion processes has been reported in physical and social environments, and corresponding models have been proposed in various areas such as electromagnetic fluids [85], ground-water solute transport [49], biology [41], finance [42], human travel [32] and predator search patterns [117].

Of the many possible models of anomalous diffusion, we shall be interested here in so-called fractional diffusion, which is a nonlocal process: to evaluate fractional diffusion at a spatial point, information involving all spatial points is needed. Recently, the analysis of such operators has received a tremendous attention: fractional diffusion has been one of the most studied topics in the past decade [38, 39, 116].

The main goal of this work is to review different techniques to approximate the solution of problems involving fractional diffusion. To make matters precise, we will consider the fractional powers of the Dirichlet Laplace operator $(-\Delta)^s$, which we will simply call the fractional Laplacian. Given $0 < s < 1$, a bounded Lipschitz domain $\Omega \subset \mathbb{R}^d$, and a function $f : \Omega \rightarrow \mathbb{R}$, we shall be concerned with finding $u : \Omega \rightarrow \mathbb{R}$ such that

$$(1.1) \quad (-\Delta)^s u = f \quad \text{in } \Omega,$$

with vanishing Dirichlet boundary conditions (understood in a suitable sense). We must immediately comment that the efficient approximation of solutions to (1.1)

Date: Draft version of August 25, 2017.

AB is supported in part by NSF grant DMS-1254618.

JPB has been partially supported by a CONICET doctoral fellowship.

RHN has been supported in part by NSF grant DMS-1411808.

EO has been supported in part by CONICYT through project FONDECYT 3160201.

AJS is supported by NSF grant DMS-1418784.

carries two essential difficulties. The first and most important one is that $(-\Delta)^s$ is *non-local*. The second feature is the lack of boundary regularity, which leads to reduced rates of convergence.

In what follows we review different definitions for the fractional Laplacian. For functions defined over \mathbb{R}^d , there is a natural way to define the fractional Laplacian as a pseudo-differential operator with symbol $|\xi|^{2s}$; namely, given a function w in the Schwartz class \mathcal{S} , set

$$(1.2) \quad (-\Delta)^s w := \mathcal{F}^{-1} (|\xi|^{2s} \mathcal{F} w),$$

where \mathcal{F} denotes the Fourier transform. The fractional Laplacian can be equivalently defined by means of the following point-wise formula (see [80, Section 1.1] and [52, Proposition 3.3])

$$(1.3) \quad (-\Delta)^s w(x) = C(d, s) \text{ p.v. } \int_{\mathbb{R}^d} \frac{w(x) - w(x')}{|x - x'|^{d+2s}} dx', \quad C(d, s) = \frac{2^{2s} s \Gamma(s + \frac{d}{2})}{\pi^{d/2} \Gamma(1 - s)},$$

where p.v. stands for the Cauchy principal value and $C(d, s)$ is a normalization constant chosen so that definitions (1.2) and (1.3) coincide. This clearly displays the non-local structure of $(-\Delta)^s w$. We remark that, in the theory of stochastic processes, expression (1.3) appears as the infinitesimal generator of a $2s$ -stable Lévy process [19].

If Ω is a bounded domain, we consider two possible definitions of the fractional Laplacian. For $u : \Omega \rightarrow \mathbb{R}$, we first extend u by zero outside Ω and next use definition (1.3). This gives the following reinterpretation of (1.1):

$$(1.4) \quad (-\Delta)^s \tilde{u} = f \quad \text{in } \Omega, \quad \tilde{u} = 0 \quad \text{in } \Omega^c = \mathbb{R}^d \setminus \Omega,$$

where the operator $(-\Delta)^s$ is understood as in (1.3) and \tilde{w} is the extension by zero to \mathbb{R}^d of a function $w : \Omega \rightarrow \mathbb{R}$ in $L^2(\Omega)$. This definition maintains the probabilistic interpretation of the fractional Laplacian defined over \mathbb{R}^d , that is, as the generator of a random walk in Ω with arbitrarily long jumps, where particles are killed upon reaching Ω^c ; see [33, Chapter 2]. The operator in (1.3) is well defined for smooth, compactly supported functions. Consequently, (1.3) can be extended by density to $\mathbb{H}^s(\Omega)$ which, for $s \in [0, 3/2)$, is defined by

$$(1.5) \quad \mathbb{H}^s(\Omega) := \{w|_{\Omega} : w \in H^s(\mathbb{R}^d), w|_{\mathbb{R}^d \setminus \Omega} = 0\}.$$

When $\partial\Omega$ is Lipschitz this space is equivalent to $\mathbb{H}^s(\Omega) = [L^2(\Omega), H_0^1(\Omega)]_s$, the real interpolation between $L^2(\Omega)$ and $H_0^1(\Omega)$ [87, 120] when $s \in [0, 1]$ and to $H^s(\Omega) \cap H_0^1(\Omega)$ when $s \in (1, 3/2)$. In what follows, we will denote by $\mathbb{H}^{-s}(\Omega)$ the dual of $\mathbb{H}^s(\Omega)$ and by $\langle \cdot, \cdot \rangle$ the duality pairing between these two spaces.

The second definition of $(-\Delta)^s$ relies on spectral theory [20]. Since $-\Delta : \mathcal{D}(-\Delta) \subset L^2(\Omega) \rightarrow L^2(\Omega)$ is an unbounded, positive and closed operator with dense domain $\mathcal{D}(-\Delta) = H_0^1(\Omega) \cap H^2(\Omega)$ and its inverse is compact, there is a countable collection of eigenpairs $\{\lambda_k, \varphi_k\}_{k \in \mathbb{N}} \subset \mathbb{R}^+ \times H_0^1(\Omega)$ such that $\{\varphi_k\}_{k \in \mathbb{N}}$ is an orthonormal basis of $L^2(\Omega)$ as well as an orthogonal basis of $H_0^1(\Omega)$. Fractional powers of the Dirichlet Laplacian can be thus defined as

$$(1.6) \quad (-\Delta)^s w := \sum_{k=1}^{\infty} \lambda_k^s w_k \varphi_k, \quad w_k := \int_{\Omega} w \varphi_k dx, \quad k \in \mathbb{N},$$

for any $w \in C_0^\infty(\Omega)$. This definition of $(-\Delta)^s$ can also be extended by density to the space $\mathbb{H}^s(\Omega)$. We also remark that, for $s \in [-1, 1]$, the space $\mathbb{H}^s(\Omega)$ can equivalently be defined by

$$\mathbb{H}^s(\Omega) = \left\{ w = \sum_{k=1}^{\infty} w_k \varphi_k : \sum_{k=1}^{\infty} \lambda_k^s w_k^2 < \infty \right\}.$$

These two definitions of the fractional Laplacian, the *integral* one involved in problem (1.4) and the *spectral* one given as in (1.6), do not coincide. In fact, as shown in [92], their difference is positive and positivity preserving, see also [36, 115]. This, in particular, implies that the boundary behavior of the solutions of (1.4) and (1.6) is quite different. According to Grubb [66, formulas (7.7)–(7.12)] the solution u of (1.4) is of the form

$$(1.7) \quad u(x) \approx \text{dist}(x, \partial\Omega)^s + v(x),$$

with v smooth; hereafter $\text{dist}(x, \partial\Omega)$ indicates the distance from $x \in \Omega$ to $\partial\Omega$. In contrast, Caffarelli and Stinga [36, Theorem 1.3] showed that solutions of (1.1), with $(-\Delta)^s$ defined as in (1.6), behave like

$$(1.8) \quad \begin{aligned} u(x) &\approx \text{dist}(x, \partial\Omega)^{2s} + v(x) & 0 < s < \frac{1}{2}; \\ u(x) &\approx \text{dist}(x, \partial\Omega) + v(x) & \frac{1}{2} < s < 1. \end{aligned}$$

The case $s = 1/2$ is exceptional; for $\Omega \subset \mathbb{R}^2$ with $\partial\Omega$ smooth, it holds that [48]

$$u(x) \approx \text{dist}(x, \partial\Omega) |\log(\text{dist}(x, \partial\Omega))| + v(x),$$

with v smooth. This lack of boundary regularity is responsible for reduced rates of convergence.

The presence of the non-integrable kernel in (1.3) is a notorious numerical difficulty that has hampered progress in the multidimensional case $d > 1$ until recently. If $\Omega = \mathbb{R}^d$, the Caffarelli-Silvestre [38] extension converts (1.1) into the following Dirichlet-to-Neumann map formulated in the cylinder $\mathcal{C} = \Omega \times (0, \infty)$

$$(1.9) \quad \text{div}(y^\alpha \nabla \mathcal{U}) = 0 \text{ in } \mathcal{C}, \quad \partial_{\nu^\alpha} \mathcal{U} = d_s f \text{ on } \mathbb{R}^d \times \{0\},$$

where $y > 0$ is the extended variable, $d_s := 2^{1-2s}\Gamma(1-s)/\Gamma(s)$ is a positive normalization constant that depends only on s , and the parameter α is defined as $\alpha := 1 - 2s \in (-1, 1)$. The relation between (1.1) and (1.9) is the following:

$$u = \mathcal{U}(\cdot, 0).$$

Cabr e and Tan [34] and Stinga and Torrea [119] have shown that a similar extension is valid for the spectral fractional Laplacian in a bounded domain Ω provided that a vanishing Dirichlet condition is appended on the lateral boundary $\partial_L \mathcal{C} = \partial\Omega \times (0, \infty)$; see also [40, 30]. Although (1.9) is a local problem, and thus amenable to PDE techniques, it is formulated in one higher dimension and exhibits a singular character as $y \downarrow 0$.

The solution to (1.1) with either definition (1.3) or (1.6) of the fractional Laplacian can be represented using Dunford-Taylor integrals [83, Chapter 4], [124, Section VIII.7]. Let us first explain the construction for definition (1.6). For $s \in (0, 1)$ and $w \in \mathbb{H}^{-s}(\Omega)$ we have

$$(1.10) \quad (-\Delta)^{-s} w = \frac{1}{2\pi i} \int_{\mathcal{C}} z^{-s} (z + \Delta)^{-1} w \, dz,$$

where \mathcal{C} is a Jordan curve oriented to have the spectrum of $-\Delta$ to its right, and z^{-s} is defined using the principal value of $\text{Log}(z)$. In addition, since the operator $-\Delta$ is positive, one can continuously deform the contour \mathcal{C} onto the negative real axis (around the branch cut) to obtain the so-called Balakrishnan formula

$$(1.11) \quad (-\Delta)^{-s} w = \frac{\sin(s\pi)}{\pi} \int_0^\infty \mu^{-s} (\mu - \Delta)^{-1} w \, d\mu;$$

see [83, eq. (4.4)] also [124, Section IX.11] and [20, Section 10.4] for a different derivation of (1.11) using semigroup theory. This formula will be the starting point

for our Dunford-Taylor approach for the spectral fractional Laplacian. These considerations, however, cannot be carried out for the the integral fractional Laplacian (1.3) since neither (1.10) or (1.11) are well defined quantities for this operator. Therefore we will, instead, multiply (1.4) by a test function w , integrate over \mathbb{R}^d and use Parseval's equality to obtain the following weak formulation of (1.4): Given $f \in \mathbb{H}^{-s}(\Omega)$ find $u \in \mathbb{H}^s(\Omega)$ such that

$$(1.12) \quad \int_{\mathbb{R}^d} |\xi|^s \mathcal{F}(\tilde{u}) |\xi|^s \overline{\mathcal{F}(\tilde{w})} d\xi = \langle f, w \rangle \quad \forall w \in \mathbb{H}^s(\Omega),$$

where \bar{z} denotes the complex conjugate of $z \in \mathbb{C}$. Using again the Fourier transform and Parseval's equality, the left hand side of the above relation can be equivalently written as (see Theorem 4.5)

$$(1.13) \quad \frac{2 \sin(s\pi)}{\pi} \int_0^\infty \mu^{1-2s} \int_{\Omega} ((-\Delta)(I - \mu^2 \Delta)^{-1} \tilde{u}(x)) w(x) dx d\mu.$$

These ideas will be the starting point of the Dunford-Taylor method for the integral fractional Laplacian (1.3).

The purpose of this paper is to present and briefly analyze three finite element methods (FEMs) to approximate (1.1). The first will consider the *spectral* definition (1.6), the second will deal with the *integral* definition (1.3), while the third approach will be able to account for both operators by means of either (1.11) or (1.13). We must immediately remark that for special domain geometries, such as when $d = 2$ and Ω is a rectangle, the use of spectral methods can be quite efficient but we do not elaborate any further as we are interested in techniques that apply to general domains. Our presentation is organized as follows: In Section 2, we present a method that hinges on the extension (1.9) and uses PDE techniques. The second method deals with the integral formulation and is discussed in Section 3. Finally, the third method is based on exponentially convergent quadrature approximations of (1.11) for the spectral Laplacian and of (1.13) for the integral Laplacian. They are discussed in Section 4.

As usual, we write $a \lesssim b$ to mean $a \leq Cb$, with a constant C that neither depends on a, b or the discretization parameters and might change at each occurrence. Moreover, $a \approx b$ indicates $a \lesssim b$ and $b \lesssim a$.

2. THE SPECTRAL FRACTIONAL LAPLACIAN

In this section we deal with the spectral definition of the fractional Laplacian (1.6) and, on the basis of (1.9), its discretization via PDE techniques as originally developed in [95]. We must immediately remark that many of the results of this section and section 4 extend to more general symmetric elliptic operators of the form $Lw = -\operatorname{div}(A\nabla w) + cw$, with $A \in C^{0,1}(\bar{\Omega}, \mathbf{GL}(\mathbb{R}^d))$ symmetric and positive definite and $0 \leq c \in C^{0,1}(\bar{\Omega}, \mathbb{R})$.

Let Ω be a convex polytopal domain. Besides the semi-infinite cylinder $\mathcal{C} = \Omega \times (0, \infty)$, we introduce the truncated cylinder $\mathcal{C}_{\mathcal{Y}} = \Omega \times (0, \mathcal{Y})$ with height \mathcal{Y} and its lateral boundary $\partial_L \mathcal{C}_{\mathcal{Y}} = \partial\Omega \times (0, \mathcal{Y})$. Since we deal with objects defined in both \mathbb{R}^d and \mathbb{R}^{d+1} , it is convenient to distinguish the extended variable y . For $\mathbf{x} \in \mathbb{R}^{d+1}$, we denote

$$\mathbf{x} = (x, y) = (x, x_{d+1}), \quad x \in \mathbb{R}^d, \quad y \in \mathbb{R}^+.$$

2.1. Extension Property. The groundbreaking extension (1.9) of Caffarelli and Silvestre [38], valid for any power $s \in (0, 1)$, is formulated in \mathbb{R}^d . Cabré and Tan [34] and Stinga and Torrea [119] realized that a similar extension holds for the spectral Laplacian over Ω bounded; see also [40, 30]. This problem reads

$$(2.1) \quad \operatorname{div}(y^\alpha \nabla \mathcal{U}) = 0 \text{ in } \mathcal{C}, \quad \mathcal{U} = 0 \text{ on } \partial_L \mathcal{C}, \quad \partial_{\nu^\alpha} \mathcal{U} = d_s f \text{ on } \Omega \times \{0\},$$

where $\alpha = 1 - 2s$ and the so-called conormal exterior derivative of \mathcal{U} at $\Omega \times \{0\}$ is

$$(2.2) \quad \partial_{\nu^\alpha} \mathcal{U} = - \lim_{y \rightarrow 0^+} y^\alpha \partial_y \mathcal{U}(\cdot, y).$$

The limit in (2.2) must be understood in the distributional sense [34, 38, 40, 119]. With this construction at hand, the fractional Laplacian and the Dirichlet-to-Neumann operator of problem (2.1) are related by

$$d_s(-\Delta)^s u = \partial_{\nu^\alpha} \mathcal{U} \quad \text{in } \Omega.$$

The operator in (2.1) is in divergence form and thus amenable to variational techniques. However, it is *nonuniformly* elliptic because the weight y^α either blows up for $-1 < \alpha < 0$ or degenerates for $0 < \alpha < 1$ as $y \downarrow 0$; the exceptional case $\alpha = 0$ corresponds to the regular harmonic extension for $s = \frac{1}{2}$ [34]. This entails dealing with weighted Lebesgue and Sobolev spaces with the weight $|y|^\alpha$ for $\alpha \in (-1, 1)$ [30, 34, 38, 40]. Such a weight belongs to the Muckenhoupt class $A_2(\mathbb{R}^{d+1})$, which is the collection of weights ω so that [54, 60, 63, 91, 122]

$$C_{2,\omega} = \sup_B \left(\int_B \omega \, d\mathbf{x} \right) \left(\int_B \omega^{-1} \, d\mathbf{x} \right) < \infty,$$

where the supremum is taken over all balls B in \mathbb{R}^{d+1} and \int_B stands for the mean value over B . The Muckenhoupt characteristic $C_{2,\omega}$ appears in all estimates involving ω .

If $D \subset \mathbb{R}^d \times \mathbb{R}^+$, we define $L^2(y^\alpha, D)$ as the Lebesgue space for the measure $y^\alpha \, d\mathbf{x}$. We also define the weighted Sobolev space

$$H^1(y^\alpha, D) = \{w \in L^2(y^\alpha, D) : |\nabla w| \in L^2(y^\alpha, D)\},$$

where ∇w is the distributional gradient of w . We equip $H^1(y^\alpha, D)$ with the norm

$$(2.3) \quad \|w\|_{H^1(y^\alpha, D)} = \left(\|w\|_{L^2(y^\alpha, D)}^2 + \|\nabla w\|_{L^2(y^\alpha, D)}^2 \right)^{\frac{1}{2}}.$$

The space $H^1(y^\alpha, D)$ is Hilbert with the norm (2.3) and $C^\infty(D) \cap H^1(y^\alpha, D)$ is dense in $H^1(y^\alpha, D)$ because $|y|^\alpha \in A_2(\mathbb{R}^{d+1})$ (cf. [63, Theorem 1], [78] and [122, Proposition 2.1.2, Corollary 2.1.6]).

To analyze problem (2.1) we define the weighted Sobolev space

$$\mathring{H}_L^1(y^\alpha, \mathcal{C}) = \{w \in H^1(y^\alpha, \mathcal{C}) : w = 0 \text{ on } \partial_L \mathcal{C}\}.$$

The following *weighted Poincaré inequality* holds [95, inequality (2.21)]

$$\|w\|_{L^2(y^\alpha, \mathcal{C})} \lesssim \|\nabla w\|_{L^2(y^\alpha, \mathcal{C})} \quad \forall w \in \mathring{H}_L^1(y^\alpha, \mathcal{C}).$$

Consequently, the seminorm on $\mathring{H}_L^1(y^\alpha, \mathcal{C})$ is equivalent to (2.3). For $w \in H^1(y^\alpha, \mathcal{C})$, $\text{tr}_\Omega w$ denotes its trace onto $\Omega \times \{0\}$ which satisfies [95, Proposition 2.5]

$$\text{tr}_\Omega \mathring{H}_L^1(y^\alpha, \mathcal{C}) = \mathbb{H}^s(\Omega), \quad \|\text{tr}_\Omega w\|_{\mathbb{H}^s(\Omega)} \leq C_{\text{tr}_\Omega} \|w\|_{\mathring{H}_L^1(y^\alpha, \mathcal{C})}.$$

The variational formulation of (2.1) reads: find $\mathcal{U} \in \mathring{H}_L^1(y^\alpha, \mathcal{C})$ such that

$$(2.4) \quad \int_{\mathcal{C}} y^\alpha \nabla \mathcal{U} \cdot \nabla w \, d\mathbf{x} = d_s \langle f, \text{tr}_\Omega w \rangle \quad \forall w \in \mathring{H}_L^1(y^\alpha, \mathcal{C}),$$

where, we recall that, $\langle \cdot, \cdot \rangle$ corresponds to the duality pairing between $\mathbb{H}^{-s}(\Omega)$ and $\mathbb{H}^s(\Omega)$. The fundamental result of Caffarelli and Silvestre [38] for $\Omega = \mathbb{R}^d$ and of Cabré and Tan [34, Proposition 2.2] and Stinga and Torrea [119, Theorem 1.1] for Ω bounded reads: given $f \in \mathbb{H}^{-s}(\Omega)$, if $u \in \mathbb{H}^s(\Omega)$ solves (1.1) and $\mathcal{U} \in \mathring{H}_L^1(y^\alpha, \mathcal{C})$ solves (2.4), then

$$(2.5) \quad u = \text{tr}_\Omega \mathcal{U}, \quad d_s(-\Delta)^s u = \partial_{\nu^\alpha} \mathcal{U} \quad \text{in } \Omega,$$

where the first equality holds in $\mathbb{H}^s(\Omega)$, whereas the second one in $\mathbb{H}^{-s}(\Omega)$.

2.2. Regularity. To study the finite element discretization of (2.4) we must understand the regularity of \mathcal{U} . We begin by recalling that if $u = \sum_{k=1}^{\infty} u_k \varphi_k$ solves (1.1), then $u_k = \lambda_k^{-s} f_k$, with f_k being the k -th Fourier coefficient of f . The unique solution \mathcal{U} of problem (1.9) thus admits the representation [95, formula (2.24)]

$$\mathcal{U}(x, y) = \sum_{k=1}^{\infty} u_k \varphi_k(x) \psi_k(y).$$

The functions ψ_k solve the 2-point boundary value problem in \mathbb{R}^+

$$\psi_k'' + \frac{\alpha}{y} \psi_k' = \lambda_k \psi_k, \quad \text{in } (0, \infty); \quad \psi_k(0) = 1, \quad \lim_{y \rightarrow \infty} \psi_k(y) = 0.$$

Thus, if $s = \frac{1}{2}$, we have that $\psi_k(y) = \exp(-\sqrt{\lambda_k} y)$ [34, Lemma 2.10]; and, if $s \in (0, 1) \setminus \{\frac{1}{2}\}$, then [40, Proposition 2.1]

$$\psi_k(y) = c_s (\sqrt{\lambda_k} y)^s K_s(\sqrt{\lambda_k} y),$$

where $c_s = 2^{1-s}/\Gamma(s)$ and K_s denotes the modified Bessel function of the second kind [2, Chap. 9.6]. Using asymptotic properties of $K_s(\zeta)$ as $\zeta \downarrow 0$ [2, Chapter 9.6], and [102, Chap. 7.8], we obtain for $y \downarrow 0$:

$$\psi_k'(y) \approx y^{-\alpha}, \quad \psi_k''(y) \approx y^{-\alpha-1}.$$

Exploiting these estimates, the following regularity results hold [95, Theorem 2.7].

Theorem 2.1 (global regularity of the α -harmonic extension). *Let $f \in \mathbb{H}^{1-s}(\Omega)$, where $\mathbb{H}^{1-s}(\Omega)$ is defined in (1.5) for $s \in (0, 1)$. Let $\mathcal{U} \in \mathring{H}_L^1(y^\alpha, \mathcal{C})$ solve (1.9) with f as data. Then, for $s \in (0, 1) \setminus \{\frac{1}{2}\}$, we have*

$$(2.6) \quad \|\Delta_x \mathcal{U}\|_{L^2(y^\alpha, \mathcal{C})}^2 + \|\partial_y \nabla_x \mathcal{U}\|_{L^2(y^\alpha, \mathcal{C})}^2 = d_s \|f\|_{\mathbb{H}^{1-s}(\Omega)}^2,$$

and

$$(2.7) \quad \|\partial_{yy} \mathcal{U}\|_{L^2(y^\beta, \mathcal{C})} \lesssim \|f\|_{L^2(\Omega)},$$

with $\beta > 2\alpha + 1$. For the special case $s = \frac{1}{2}$, we obtain

$$\|\mathcal{U}\|_{H^2(\mathcal{C})} \lesssim \|f\|_{\mathbb{H}^{1/2}(\Omega)}.$$

Comparing (2.6) and (2.7), we realize that the regularity of \mathcal{U} is much worse in the extended $d+1$ direction. Since Ω is convex, the following elliptic regularity estimate is valid [65]:

$$(2.8) \quad \|w\|_{H^2(\Omega)} \lesssim \|\Delta_x w\|_{L^2(\Omega)} \quad \forall w \in H^2(\Omega) \cap H_0^1(\Omega).$$

This, combined with (2.6), yields the following estimate for the Hessian $D_x^2 \mathcal{U}$ in the variable $x \in \Omega$:

$$\|D_x^2 \mathcal{U}\|_{L^2(y^\alpha, \mathcal{C})} \lesssim \|f\|_{\mathbb{H}^{1-s}(\Omega)}.$$

Further regularity estimates in Hölder and Sobolev norms are derived in [36]. However, we do not need them for what follows.

2.3. Truncation. Since \mathcal{C} is unbounded, problem (2.1) cannot be approximated with standard finite element techniques. However, since the solution \mathcal{U} of problem (2.1) decays exponentially in y [95, Proposition 3.1], by truncating \mathcal{C} to $\mathcal{C}_\mathcal{Y} := \Omega \times (0, \mathcal{Y})$ and setting a homogeneous Dirichlet condition on $y = \mathcal{Y}$, we only incur in an exponentially small error in terms of \mathcal{Y} [95, Theorem 3.5]. If

$$\mathring{H}_L^1(y^\alpha, \mathcal{C}_\mathcal{Y}) := \{w \in H^1(y^\alpha, \mathcal{C}_\mathcal{Y}) : w = 0 \text{ on } \Gamma_D\},$$

where $\Gamma_D = \partial_L \mathcal{C}_\gamma \cup \Omega \times \{\gamma\}$ is the Dirichlet boundary, then the aforementioned problem reads:

$$(2.9) \quad \mathcal{V} \in \mathring{H}_L^1(y^\alpha, \mathcal{C}_\gamma) : \int_{\mathcal{C}_\gamma} y^\alpha \nabla \mathcal{V} \cdot \nabla w \, d\mathbf{x} = d_s \langle f, \text{tr}_\Omega w \rangle \quad \forall w \in \mathring{H}_L^1(y^\alpha, \mathcal{C}_\gamma).$$

The following exponential decay rate is proved in [95, Theorem 3.5].

Lemma 2.2 (truncation error). *If \mathcal{U} and \mathcal{V} denote the solutions of (2.4) and (2.9), respectively, then*

$$\|\nabla(\mathcal{U} - \mathcal{V})\|_{L^2(y^\alpha, \mathcal{C})} \lesssim e^{-\sqrt{\lambda_1} \gamma/4} \|f\|_{\mathbb{H}^{-s}(\Omega)}$$

is valid, where λ_1 denotes the first eigenvalue of the Dirichlet Laplace operator and $\gamma \geq 1$ is the truncation parameter.

2.4. FEM: A Priori Error Analysis. The first numerical work that exploits the groundbreaking identity (2.5), designs and analyzes a FEM for (2.1) is [95]; see also [96, 103]. We briefly review now the main a priori results of [95].

We introduce a conforming and shape regular mesh $\mathcal{T} = \{K\}$ of Ω , where $K \subset \mathbb{R}^d$ is an element that is isoparametrically equivalent either to the unit cube $[0, 1]^d$ or the unit simplex in \mathbb{R}^d . Over this mesh we construct the finite element space

$$(2.10) \quad \mathbb{U}(\mathcal{T}) = \{W \in C^0(\bar{\Omega}) : W|_K \in \mathcal{P}_1(K) \, \forall K \in \mathcal{T}, W|_{\partial\Omega} = 0\}.$$

The set $\mathcal{P}_1(K)$ is either the space $\mathbb{P}_1(K)$ of polynomials of total degree at most 1, when K is a simplex, or the space of polynomials $\mathbb{Q}_1(K)$ of degree not larger than 1 in each variable provided K is a d -rectangle.

To triangulate the truncated cylinder \mathcal{C}_γ we consider a partition \mathcal{I}_γ of the interval $[0, \gamma]$ with nodes y_m , $m = 0, \dots, M$, and construct a mesh \mathcal{T}_γ of \mathcal{C}_γ as the tensor product of \mathcal{T} and \mathcal{I}_γ . The set of all triangulations of \mathcal{C}_γ obtained with this procedure is \mathbb{T} .

For $\mathcal{T}_\gamma \in \mathbb{T}$, we define the finite element space

$$(2.11) \quad \mathbb{V}(\mathcal{T}_\gamma) = \{W \in C^0(\bar{\mathcal{C}}_\gamma) : W|_T \in \mathcal{P}_1(K) \otimes \mathbb{P}_1(I) \, \forall T \in \mathcal{T}_\gamma, W|_{\Gamma_D} = 0\},$$

where $T = K \times I$ and observe that $\mathbb{U}(\mathcal{T}) = \text{tr}_\Omega \mathbb{V}(\mathcal{T}_\gamma)$. Note that $\#\mathcal{T}_\gamma = M \#\mathcal{T}$, and that $\#\mathcal{T} \approx M^d$ implies $\#\mathcal{T}_\gamma \approx M^{d+1}$.

The Galerkin approximation of (2.9) is the function $V \in \mathbb{V}(\mathcal{T}_\gamma)$ such that

$$(2.12) \quad \int_{\mathcal{C}_\gamma} y^\alpha \nabla V \cdot \nabla W \, d\mathbf{x} = d_s \langle f, \text{tr}_\Omega W \rangle \quad \forall W \in \mathbb{V}(\mathcal{T}_\gamma).$$

Existence and uniqueness of V immediately follows from $\mathbb{V}(\mathcal{T}_\gamma) \subset \mathring{H}_L^1(y^\alpha, \mathcal{C}_\gamma)$ and the Lax-Milgram Lemma. It is trivial also to obtain a best approximation result *à la* Céa, namely

$$(2.13) \quad \|\nabla(\mathcal{V} - V)\|_{L^2(y^\alpha, \mathcal{C}_\gamma)} = \inf_{W \in \mathbb{V}(\mathcal{T}_\gamma)} \|\nabla(\mathcal{V} - W)\|_{L^2(y^\alpha, \mathcal{C}_\gamma)}.$$

This reduces the numerical analysis of (2.12) to a question in approximation theory, which in turn can be answered by the study of piecewise polynomial interpolation in Muckenhoupt weighted Sobolev spaces; see [95, 98]. Exploiting the Cartesian structure of the mesh \mathcal{T}_γ we are able to extend the anisotropic estimates of Durán and Lombardi [55] to our setting [95, Theorems 4.6–4.9], [98]. The following error estimates separate in each direction.

Proposition 2.3 (anisotropic interpolation estimates). *There exists a quasi interpolation operator $\Pi_{\mathcal{T}_y} : L^1(\mathcal{C}_y) \rightarrow \mathbb{V}(\mathcal{T}_y)$ that satisfies the following anisotropic error estimates for all $j = 1, \dots, d+1$ and all $T = K \times I \in \mathcal{T}_y$*

$$\begin{aligned} \|w - \Pi_{\mathcal{T}_y} w\|_{L^2(y^\alpha, T)} &\lesssim h_K \|\nabla_x w\|_{L^2(y^\alpha, S_T)} + h_I \|\partial_y w\|_{L^2(y^\alpha, S_T)}, \\ \|\partial_{x_j}(w - \Pi_{\mathcal{T}_y} w)\|_{L^2(y^\alpha, T)} &\lesssim h_K \|\nabla_x \partial_{x_j} w\|_{L^2(y^\alpha, S_T)} + h_I \|\partial_y \partial_{x_j} w\|_{L^2(y^\alpha, S_T)}, \end{aligned}$$

where S_T stands for the patch of elements of \mathcal{T}_y that intersect T , $h_K = |K|^{1/d}$ and $h_I = |I|$.

As a first application of Proposition 2.3 we consider a quasiuniform mesh \mathcal{T}_y of size h and set $w = \mathcal{U}$. We estimate $\mathcal{U} - \Pi_{\mathcal{T}_y} \mathcal{U}$ for $y \geq 2h$ as follows:

$$\int_{2h}^{\mathcal{Y}} y^\alpha \|\partial_y(\mathcal{U} - \Pi_{\mathcal{T}_y} \mathcal{U})\|_{L^2(\Omega)}^2 dy \lesssim h^2 \int_h^{\mathcal{Y}} y^\alpha \left(\|\partial_{yy} \mathcal{U}\|_{L^2(\Omega)}^2 + \|\nabla_x \partial_y \mathcal{U}\|_{L^2(\Omega)}^2 \right) dy.$$

For the first term we resort to (2.7), and recall that $\beta > 2\alpha + 1 > \alpha$, to deduce

$$\begin{aligned} h^2 \int_h^{\mathcal{Y}} y^\alpha \|\partial_{yy} \mathcal{U}\|_{L^2(\Omega)}^2 dy &\leq h^2 \sup_{h \leq y \leq \mathcal{Y}} y^{\alpha-\beta} \int_0^y y^\beta \|\partial_{yy} \mathcal{U}\|_{L^2(\Omega)}^2 dy \\ &\leq h^{2+\alpha-\beta} \|f\|_{L^2(\Omega)}^2. \end{aligned}$$

For the second term we use (2.6) instead to arrive at

$$h^2 \int_h^{\mathcal{Y}} y^\alpha \|\nabla_x \partial_y \mathcal{U}\|_{L^2(\Omega)}^2 dy \leq h^2 \int_0^{\mathcal{Y}} y^\alpha \|\nabla_x \partial_y \mathcal{U}\|_{L^2(\Omega)}^2 dy \lesssim h^2 \|f\|_{\mathbb{H}^{1-s}(\Omega)}^2.$$

Combining the two estimates we derive the interpolation estimate

$$\int_{2h}^{\mathcal{Y}} y^\alpha \|\partial_y(\mathcal{U} - \Pi_{\mathcal{T}_y} \mathcal{U})\|_{L^2(\Omega)}^2 dy \lesssim h^{2+\alpha-\beta} \|f\|_{\mathbb{H}^{1-s}(\Omega)}^2.$$

To bound $\mathcal{U} - \Pi_{\mathcal{T}_y} \mathcal{U}$ for $0 \leq y \leq 2h$ we invoke the stability of the operator $\Pi_{\mathcal{T}_y}$ [95, Theorems 4.7 and 4.8] and the asymptotic behavior of \mathcal{U} : $\partial_y \mathcal{U} \approx y^{-\alpha}$ as $y \approx 0^+$. This yields

$$\int_0^{2h} y^\alpha \|\partial_y(\mathcal{U} - \Pi_{\mathcal{T}_y} \mathcal{U})\|_{L^2(\Omega)}^2 dy \lesssim h^{1-\alpha} \|f\|_{\mathbb{H}^{1-s}(\Omega)}^2.$$

The previous estimates, combined with the fact that $2 + \alpha + \beta < 1 - \alpha$, allow us to derive that

$$\int_0^{\mathcal{Y}} y^\alpha \|\partial_y(\mathcal{U} - \Pi_{\mathcal{T}_y} \mathcal{U})\|_{L^2(\Omega)}^2 dy \lesssim h^{2+\alpha-\beta} \|f\|_{\mathbb{H}^{1-s}(\Omega)}^2,$$

which is quasi-optimal in terms of regularity because $2 + \alpha - \beta = 2(s - \epsilon)$ and $\partial_y \mathcal{U} \approx y^{-\alpha}$ formally implies $\mathcal{U} \in H^{s-\epsilon}(y^\alpha, \mathcal{C})$ for any $\epsilon > 0$; however this estimate exhibits a suboptimal rate. To restore a quasi-optimal rate, we must compensate the behavior of $\partial_{yy} \mathcal{U}$ by a *graded* mesh in the extended direction, which is allowed by Proposition 2.3. Therefore, we construct a mesh \mathcal{I}_y with nodes

$$(2.14) \quad y_m = m^\gamma M^{-\gamma} \mathcal{Y}, \quad m = 0, \dots, M,$$

where $\gamma > 3/(1 - \alpha) = 3/(2s)$. Combining (2.13) with Proposition 2.3 we obtain estimates in terms of degrees of freedom [95, Theorem 5.4 and Corollary 7.11].

Theorem 2.4 (a priori error estimate). *Let $\mathcal{T}_y = \mathcal{T} \times \mathcal{I}_y \in \mathbb{T}$ with \mathcal{I}_y satisfying (2.14), and let $\mathbb{V}(\mathcal{T}_y)$ be defined by (2.11). If $V \in \mathbb{V}(\mathcal{T}_y)$ solves (2.12), we have*

$$(2.15) \quad \|\nabla(\mathcal{U} - V)\|_{L^2(y^\alpha, \mathcal{C})} \lesssim |\log(\#\mathcal{T}_y)|^s (\#\mathcal{T}_y)^{-1/(d+1)} \|f\|_{\mathbb{H}^{1-s}(\Omega)},$$

where $\mathcal{Y} \approx \log(\#\mathcal{T}_y)$. Alternatively, if u denotes the solution of (1.1), then

$$\|u - U\|_{\mathbb{H}^s(\Omega)} \lesssim |\log(\#\mathcal{T}_y)|^s (\#\mathcal{T}_y)^{-1/(d+1)} \|f\|_{\mathbb{H}^{1-s}(\Omega)},$$

where $U = \text{tr}_\Omega V$.

Remark 2.5 (domain and data regularity). The estimates of Theorem 2.4 hold only if $f \in \mathbb{H}^{1-s}(\Omega)$ and the domain Ω is such that (2.8) is valid.

Remark 2.6 (quasi-uniform meshes). Let $V \in \mathbb{V}(\mathcal{T}_\mathcal{Y})$ solve (2.12) over a quasi-uniform mesh $\mathcal{T}_\mathcal{Y}$ of $\mathcal{C}_\mathcal{Y}$ of size h . Combining (2.13) with the preceding discussion we obtain for $\mathcal{Y} \approx |\log h|$ and all $\varepsilon > 0$

$$\|\nabla(\mathcal{U} - V)\|_{L^2(y^\alpha, \mathcal{C}_\mathcal{Y})} \lesssim h^{s-\varepsilon} \|f\|_{\mathbb{H}^{1-s}(\Omega)},$$

where the hidden constant blows up if $\varepsilon \downarrow 0$.

Remark 2.7 (case $s = \frac{1}{2}$). If $s = \frac{1}{2}$, we obtain the optimal estimate

$$\|\nabla(\mathcal{U} - V)\|_{L^2(\mathcal{C}_\mathcal{Y})} \lesssim h \|f\|_{\mathbb{H}^{1/2}(\Omega)}.$$

Remark 2.8 (complexity). Except for a logarithmic factor, the error estimates of Theorem 2.4 decay with a rate $(\#\mathcal{T}_\mathcal{Y})^{-1/(d+1)}$, where d is the dimension of Ω . For $f \in \mathbb{H}^{1-s}(\Omega)$, the error estimate (2.15) is optimal in terms of approximation. However, in terms of error versus work, the FEM (2.12) is sub-optimal as a method to compute in Ω . Recently, approaches based on hp -methods [18, 88] and sparse grids [18] have been proposed. These approaches yield methods with, up to logarithmic factors, $\mathcal{O}(\#\mathcal{T})$ degrees of freedom and realize the optimal asymptotic convergence rate $(\#\mathcal{T})^{-1/d}$, again, up to logarithmic factors; see Section 2.7 for a detailed discussion.

2.5. Numerical Experiments. We present two numerical examples for $d = 2$ computed within the `deal.II` library [16, 17] using graded meshes. Integrals are evaluated with a tensor product Gaussian quadrature with 15 nodes in each coordinate direction and linear systems are solved using CG with ILU preconditioner and the exit criterion being that the ℓ^2 -norm of the residual is less than 10^{-12} .

2.5.1. Square Domain. Let $\Omega = (0, 1)^2$. Then

$$\varphi_{m,n}(x_1, x_2) = \sin(m\pi x_1) \sin(n\pi x_2), \quad \lambda_{m,n} = \pi^2 (m^2 + n^2), \quad m, n \in \mathbb{N}.$$

If $f(x_1, x_2) = (2\pi^2)^s \sin(\pi x_1) \sin(\pi x_2)$, then $u(x_1, x_2) = \sin(\pi x_1) \sin(\pi x_2)$, by (1.6) and $\mathcal{U}(x_1, x_2, y) = 2^{1-s/2} \pi^s \Gamma(s)^{-1} \sin(\pi x_1) \sin(\pi x_2) y^s K_s(\sqrt{2}\pi y)$ [95, (2.24)].

We construct a sequence of meshes $\{\mathcal{T}_{\mathcal{Y}_k}\}_{k \geq 1}$, where \mathcal{T} is obtained by uniform refinement and $\mathcal{I}_{\mathcal{Y}_k}$ is given by (2.14) with parameter $\gamma = 3/(1-\alpha) + 1/10$. On the basis of Theorem 2.4, the truncation parameter \mathcal{Y}_k is chosen to be

$$\mathcal{Y}_k = \log(\#\mathcal{T}_{\mathcal{Y}_{k-1}})^{-1/3} + 1.$$

With this type of meshes,

$$\|u - \text{tr}_\Omega V_k\|_{\mathbb{H}^s(\Omega)} \lesssim \|\nabla(\mathcal{U} - V_k)\|_{L^2(y^\alpha, \mathcal{C})} \lesssim |\log(\#\mathcal{T}_{\mathcal{Y}_k})|^s \cdot (\#\mathcal{T}_{\mathcal{Y}_k})^{-1/3},$$

which is near-optimal in \mathcal{U} but suboptimal in u , since we should expect (see [31])

$$\|u - \text{tr}_\Omega V_k\|_{\mathbb{H}^s(\Omega)} \lesssim h^{2-s} \lesssim (\#\mathcal{T}_{\mathcal{Y}_k})^{-(2-s)/3}.$$

Figure 1 shows the rates of convergence for $s = 0.2$ and $s = 0.8$ respectively. In both cases, we obtain the rate given by Theorem 2.4.




FIGURE 1. Computational rate of convergence for a square and graded meshes. The left panel shows the rate for $s = 0.2$ and the right one for $s = 0.8$. The experimental rate of convergence is $(\#\mathcal{T}_\gamma)^{-1/3}$, in agreement with Theorem 2.4.




FIGURE 2. Computational rate of convergence for a circle with graded meshes. The left panel shows the rate for $s = 0.3$ and the right one for $s = 0.7$. The experimental rate of convergence is $(\#\mathcal{T}_\gamma)^{-1/3}$, in agreement with Theorem 2.4.

2.5.2. *Circular Domain.* If $\Omega = \{x \in \mathbb{R}^2 : |x| < 1\}$, then

$$\varphi_{m,n}(r, \theta) = J_m(j_{m,n}r) (A_{m,n} \cos(m\theta) + B_{m,n} \sin(m\theta)), \quad \lambda_{m,n} = j_{m,n}^2,$$

where J_m is the m -th Bessel function of the first kind; $j_{m,n}$ is the n -th zero of J_m and $A_{m,n}, B_{m,n}$ are normalization constants to ensure $\|\varphi_{m,n}\|_{L^2(\Omega)} = 1$.

If $f = (\lambda_{1,1})^s \varphi_{1,1}$, then (1.6) and [95, (2.24)] show that $u = \varphi_{1,1}$ and

$$\mathcal{U}(r, \theta, y) = 2^{1-s} \Gamma(s)^{-1} (\lambda_{1,1})^{s/2} \varphi_{1,1}(r, \theta) y^s K_s(\sqrt{2\pi}y).$$

We construct a sequence of meshes $\{\mathcal{T}_{\gamma_k}\}_{k \geq 1}$ as in §2.5.1. With these meshes

$$\|\mathcal{U} - V_k\|_{\hat{H}_L^1(y^\alpha, \mathcal{C})} \lesssim |\log(\#\mathcal{T}_{\gamma_k})|^s (\#\mathcal{T}_{\gamma_k})^{-1/3},$$

which is near-optimal. Figure 2 shows the errors of $\|\mathcal{U} - V_k\|_{H^1(y^\alpha, \mathcal{C}_{\gamma_k})}$ for $s = 0.3$ and $s = 0.7$. The results, again, are in agreement with Theorem 2.4.

2.6. FEM: A Posteriori Error Analysis. A posteriori error estimation and adaptive finite element methods (AFEMs) have been the subject of intense research since the late 1970's because they yield optimal performance in situations where classical FEM cannot. The a priori theory for (2.12) requires $f \in \mathbb{H}^{1-s}(\Omega)$ and Ω convex for (2.8) to be valid; see Remark 2.5. If either of these does not hold, then \mathcal{U} may have singularities in the x -variables and Theorem 2.4 may not apply: a quasi-uniform refinement of Ω would not result in an efficient solution technique. An adaptive loop driven by an a posteriori error estimator is essential to recover optimal rates of convergence. In what follows we explore this.

We first observe that we cannot rely on residual error estimators. In fact, they hinge on the strong form of the local residual to measure the error. Let $T \in \mathcal{T}_y$ and ν be the unit outer normal to T . Integration by parts yields

$$\int_T y^\alpha \nabla V \cdot \nabla W \, d\mathbf{x} = \int_{\partial T} y^\alpha W \nabla V \cdot \nu \, d\sigma - \int_T \operatorname{div}(y^\alpha \nabla V) W \, d\mathbf{x}.$$

Since $\alpha \in (-1, 1)$ the boundary integral is meaningless for $y = 0$. Even the very first step in the derivation of a residual a posteriori error estimator fails! There is nothing left to do but to consider a different type of estimator.

2.6.1. Local Problems over Cylindrical Stars. Inspired by [13, 90], we deal with the anisotropy of the mesh in the extended variable y and the coefficient y^α by considering local problems on cylindrical stars. The solutions of these local problems allow us to define an anisotropic a posteriori error estimator which, under certain assumptions, is equivalent to the error up to data oscillation terms.

Given a node \mathbf{v} on the mesh \mathcal{T}_y , we exploit the tensor product structure of \mathcal{T}_y , and we write $\mathbf{v} = (\mathbf{v}, \bar{\mathbf{w}})$ where \mathbf{v} and $\bar{\mathbf{w}}$ are nodes on the meshes \mathcal{T} and \mathcal{L}_y respectively. For $K \in \mathcal{T}$, we denote by $\mathcal{N}(K)$ and $\mathring{\mathcal{N}}(K)$ the set of nodes and interior nodes of K , respectively. We set

$$\mathcal{N}(\mathcal{T}) = \bigcup_{K \in \mathcal{T}} \mathcal{N}(K), \quad \mathring{\mathcal{N}}(\mathcal{T}) = \bigcup_{K \in \mathcal{T}} \mathring{\mathcal{N}}(K).$$

The *star* around \mathbf{v} is $S_{\mathbf{v}} = \bigcup_{K \ni \mathbf{v}} K \subset \Omega$, and the *cylindrical star* around \mathbf{v} is

$$\mathcal{C}_{\mathbf{v}} := \bigcup \{T \in \mathcal{T}_y : T = K \times I, K \ni \mathbf{v}\} = S_{\mathbf{v}} \times (0, \mathcal{I}) \subset \mathcal{C}_y.$$

For each node $\mathbf{v} \in \mathcal{N}(\mathcal{T})$ we define the local space

$$\mathbb{W}(\mathcal{C}_{\mathbf{v}}) = \{w \in H^1(y^\alpha, \mathcal{C}_{\mathbf{v}}) : w = 0 \text{ on } \partial \mathcal{C}_{\mathbf{v}} \setminus \Omega \times \{0\}\},$$

and the (ideal) estimator $\eta_{\mathbf{v}} \in \mathbb{W}(\mathcal{C}_{\mathbf{v}})$ to be the solution of

$$\int_{\mathcal{C}_{\mathbf{v}}} y^\alpha \nabla \eta_{\mathbf{v}} \cdot \nabla w \, d\mathbf{x} = d_s \langle f, \operatorname{tr}_\Omega w \rangle - \int_{\mathcal{C}_{\mathbf{v}}} y^\alpha \nabla V \cdot \nabla w \, d\mathbf{x} \quad \forall w \in \mathbb{W}(\mathcal{C}_{\mathbf{v}}).$$

We finally define the local indicators $\mathcal{E}_{\mathbf{v}}$ and global error estimators $\mathcal{E}_{\mathcal{T}_y}$, as follows:

$$(2.16) \quad \mathcal{E}_{\mathbf{v}} = \|\nabla \eta_{\mathbf{v}}\|_{L^2(y^\alpha, \mathcal{C}_{\mathbf{v}})}, \quad \mathcal{E}_{\mathcal{T}_y}^2 = \sum_{\mathbf{v} \in \mathcal{N}(\mathcal{T})} \mathcal{E}_{\mathbf{v}}^2.$$

We have the following key properties [43, Proposition 5.14].

Proposition 2.9 (a posteriori error estimates). *Let $\mathcal{V} \in \mathring{H}_L^1(y^\alpha, \mathcal{C}_y)$ and $V \in \mathbb{V}(\mathcal{T}_y)$ solve (2.9) and (2.12) respectively. Then, the estimator defined in (2.16) satisfies the global bound*

$$\|\nabla(\mathcal{V} - V)\|_{L^2(y^\alpha, \mathcal{C}_y)} \lesssim \mathcal{E}_{\mathcal{T}_y},$$

and the local bound with constant 1 for any $\mathbf{v} \in \mathcal{N}(\mathcal{T})$

$$\mathcal{E}_{\mathbf{v}} \leq \|\nabla(\mathcal{V} - V)\|_{L^2(y^\alpha, \mathcal{C}_{\mathbf{v}})}.$$

These estimates provide the best scenario for a posteriori error analysis but they are not practical because the local space $\mathbb{W}(\mathcal{C}_{\mathbf{v}})$ is infinite dimensional. We further discretize $\mathbb{W}(\mathcal{C}_{\mathbf{v}})$ with continuous piecewise polynomials of degree > 1 as follows. If K is a quadrilateral, we use polynomials of degree ≤ 2 in each variable. If K is a simplex, we employ polynomials of total degree ≤ 2 augmented by a local cubic bubble function. We tensorize these spaces with continuous piecewise quadratics in the extended variable. We next construct discrete subspaces of the local space $\mathbb{W}(\mathcal{C}_{\mathbf{v}})$ and corresponding discrete estimators instead of (2.16). Under suitable assumptions, Proposition 2.9 extends to this case [43, Section 5.4].

2.6.2. *Numerical Experiment.* We illustrate the performance of a practical version of the a posteriori error estimator (2.16). We use an adaptive loop

SOLVE \rightarrow ESTIMATE \rightarrow MARK \rightarrow REFINES

with Dörfler marking. We generate a new mesh \mathcal{T}' by bisecting all the elements $K \in \mathcal{T}$ contained in the marked set \mathcal{M} based on newest-vertex bisection method [99, 100]. We choose the truncation parameter as $\mathcal{Y} = 1 + \frac{1}{3} \log(\#\mathcal{T}')$ [95, Remark 5.5]. We set $M \approx (\#\mathcal{T}')^{1/d}$ and construct $\mathcal{I}'_{\mathcal{Y}}$ by the rule (2.14). The new mesh $\mathcal{T}'_{\mathcal{Y}} = \text{REFINE}(\mathcal{M})$ is obtained as the tensor product of \mathcal{T}' and $\mathcal{I}'_{\mathcal{Y}}$.

We consider the data $\Omega = (0, 1)^2$ and $f \equiv 1$, which is incompatible for $s < 1/2$ because f does not have a vanishing trace whence $f \notin \mathbb{H}^{1-s}(\Omega)$; see also [95, Section 6.3]. Therefore, we cannot expect an optimal rate for quasi-uniform meshes \mathcal{T} in Ω according to Theorem 2.4. Figure 3 shows that adaptive mesh refinement guided by AFEM restores an optimal decay rate for $s < 1/2$.




FIGURE 3. Computational rate of convergence of AFEM for $d = 2$ and $s = 0.2, 0.4, 0.6$ and $s = 0.8$. The left panel shows the error vs. the number of degrees of freedom, the right one the total error indicator. We recover the optimal rate $\#\mathcal{T}_{\mathcal{Y}}^{-1/3}$. For $s < \frac{1}{2}$, the right hand side $f \equiv 1 \notin \mathbb{H}^{1-s}(\Omega)$ and a quasiuniform mesh in Ω does not deliver an optimal rate of convergence [95, Section 6.3].

2.7. **Extensions and Applications.** We conclude the discussion of this approach by mentioning several extension and applications:

- *Efficient solvers:* Finding the solution to (2.9) entails solving a large linear system with a sparse matrix. In [44] the construction of efficient multilevel techniques for the solution of this problem was addressed. It was shown that multilevel techniques with line smoothers over vertical lines in the extended direction perform almost optimally; i.e., the contraction factor depends linearly on the number of levels, and thus logarithmically on the problem size.
- *Time dependent problems:* In [97] the time dependent problem

$$(2.17) \quad \partial_t^\gamma u + (-\Delta)^s u = f \quad \Omega \times (0, T), \quad u(\cdot, 0) = u_0$$

was examined. Here ∂_t^γ denotes the so-called Caputo derivative of order $\gamma \in (0, 1)$, which is defined as follows [112]:

$$(2.18) \quad \partial_t^\gamma w(x, t) := \frac{1}{\Gamma(1-\gamma)} \int_0^t \frac{1}{(t-r)^\gamma} \partial_r w(x, r) dr \quad t > 0, x \in \Omega.$$

It turns out that the solution of this problem is always singular as $t \downarrow 0$ provided the initial condition $u_0 \neq 0$. In fact, taking $f = 0$ and representing u in terms of the Mittag-Leffler function [64] reveals that

$$u(x, t) = \left(1 - \frac{t^\gamma}{\Gamma(1+\gamma)} (-\Delta)^s + \mathcal{O}(t^{2\gamma}) \right) u_0(x) \quad x \in \Omega.$$

These heuristics led to the following regularity results shown in [97]

$$(2.19) \quad \partial_t u \in L \log L(0, T; \mathbb{H}^{-s}(\Omega)), \quad \partial_{tt}^2 u \in L^2(t^\sigma; 0, T; \mathbb{H}^{-s}(\Omega)), \quad \sigma > 3 - 2\gamma,$$

where $L \log L(0, T)$ denotes the Orlicz space of functions w such that $|w| \log |w| \in L^1(0, T)$; see [76]. Using the extension property, problem (2.17) reduces to a quasi-stationary elliptic equation with a dynamic boundary condition. Rates of convergence for fully discrete schemes were derived in [97], which are consistent with the regularity (2.19); this issue has been largely ignored in the literature.

The extension of these results to a space-time fractional wave equation, i.e., $\gamma \in (1, 2]$ is currently under investigation [106].

- *Nonlinear problems:* Elliptic and parabolic obstacle problems with the spectral fractional Laplacian were considered in [94] and [105], respectively. Rates of convergence for their FEM approximation were derived, and this required a careful combination of Sobolev regularity, as in Theorem 2.1, and Hölder regularity of the solution [37, 35]. In addition, a positivity preserving interpolant that is stable in anisotropic meshes and weighted norms had to be constructed.
- *PDE constrained optimization and optimal control:* Optimal control problems where the state equation is given by either a stationary or parabolic equation with a spectral fractional Laplacian were studied in [9] and [12], respectively. Existence and uniqueness of optimal pairs was obtained, as well as their regularity. In both cases fully discrete schemes were designed and their convergence shown. The results of [9] were later improved upon considering piecewise linear approximation of the optimal control variable [104] and adaptive algorithms [10]. Sparse optimal control for the spectral fractional Laplacian was studied in [107]. Finally, reference [11] presents the design and analysis of an approximation scheme for an optimal control problem where the control variable corresponds to the order of the fractional operator [118].
- *Near optimal complexity:* According to Remark 2.8, the FEM (2.12) on anisotropic meshes with radical grading (2.14) is suboptimal as a method to compute in Ω . In fact, up to logarithmic factors, the total number of degrees of freedom involved in the method behaves like $\mathcal{O}(\#\mathcal{T}^{1+1/d})$. This deficiency has been recently cured by means of two approaches. First, in [18, 88] a full tensor product approach of a hp -FEM on the extended variable y with a P_1 -FEM in Ω yield, up to logarithmic factors, the error estimate

$$(2.20) \quad \|u - U\|_{\mathbb{H}^s(\Omega)} \lesssim (\#\mathcal{T})^{-1/d} \|f\|_{\mathbb{H}^{1-s}(\Omega)}.$$

This scheme exploits the analyticity properties of $\mathcal{U}(\cdot, y)$ in y ; see [18, Theorem 4.7]. The total number of degrees of freedom of the method behaves, and again up to logarithmic factors, like $\mathcal{O}(\#\mathcal{T})$ [18, Theorem 5.14] and thus, it is optimal as a method to compute in Ω . We must immediately remark that this solution technique *circumvents the fact that an extra dimension was incorporated to the resolution of the spectral fractional Laplacian*. Second, in [18] a novel sparse tensor product P_1 -FEM has been proposed and analyzed. The total number of degrees of freedom involved in this scheme behaves as $\mathcal{O}(\#\mathcal{T})$ and realizes the optimal error estimate (2.20).

- *Non convex domains:* In two dimensions, and by suitably grading the mesh towards the reentrant corners of the domain, the two schemes of [18] that we previously commented on yield optimal error estimates also for nonconvex domains $\Omega \subset \mathbb{R}^2$. These results rely on the weighted analytic regularity of \mathcal{U} in the extended variable y and the regularity of \mathcal{U} in Kondrat'ev spaces on Ω ; see [18, Theorem 5.5].
- *hp-methods in \mathbf{x} :* In the case of a smooth domain and a smooth, but not necessarily compatible f , [18] has also developed an hp -method over the cylinder \mathcal{C}_γ .

In this case, the results show (see [18, Theorems 7.3 and 7.7]) that *exponential* rates of convergence with respect to the number of degrees of freedom can be achieved. This is the first and only scheme, at the time of this writing, that circumvents the compatibility condition $f \in \mathbb{H}^{1-s}(\Omega)$.

- *Diagonalization*: Finally, in [18, Section 6], a diagonalization procedure was introduced which reduces this approach to the solution of M independent and embarrassingly parallelizable elliptic problems. Thus, for instance, for the *hp*-version of this method, one only needs to solve $\log(\#\mathcal{T})$ problems of size $\#\mathcal{T}$.

3. THE INTEGRAL FRACTIONAL LAPLACIAN

Here we consider the discretization of the integral definition of the fractional Laplacian (1.4). In view of the definition (1.5) of the space $\mathbb{H}^s(\Omega)$, and the fractional Poincaré inequality

$$\|w\|_{L^2(\Omega)} \lesssim |w|_{H^s(\mathbb{R}^d)} \quad \forall w \in \mathbb{H}^s(\Omega),$$

we may furnish $\mathbb{H}^s(\Omega)$ with the $H^s(\mathbb{R}^d)$ -seminorm. We also define the bilinear form $[[\cdot, \cdot]] : \mathbb{H}^s(\Omega) \times \mathbb{H}^s(\Omega) \rightarrow \mathbb{R}$,

$$(3.1) \quad [[u, w]] := \frac{C(d, s)}{2} \iint_Q \frac{(u(x) - u(x'))(w(x) - w(x'))}{|x - x'|^{d+2s}} dx' dx,$$

where $Q = (\Omega \times \mathbb{R}^d) \cup (\mathbb{R}^d \times \Omega)$ and $C(d, s)$ was defined in (1.3). We denote by $||| \cdot |||$ the norm that $[[\cdot, \cdot]]$ induces, which is just a multiple of the $H^s(\mathbb{R}^d)$ -seminorm. The weak formulation of (1.4) is obtained upon multiplying (1.3) by a test function $w \in \mathbb{H}^s(\Omega)$, integrating over $x \in \Omega$ and exploiting symmetry to make the difference $w(x) - w(x')$ appear. With the functional setting we have just described at hand, this problem is formulated as follows: find $u \in \mathbb{H}^s(\Omega)$ such that

$$(3.2) \quad [[u, w]] = \langle f, w \rangle \quad \forall w \in \mathbb{H}^s(\Omega),$$

where, we recall that $\langle \cdot, \cdot \rangle$ corresponds to the duality pairing between $\mathbb{H}^{-s}(\Omega)$ and $\mathbb{H}^s(\Omega)$. Applying the Lax-Milgram lemma immediately yields well-posedness of (3.2).

Although the energy norm $||| \cdot |||$ involves integration on $\Omega \times \mathbb{R}^d$, this norm can be localized. In fact, due to Hardy's inequality [45, 56], the following equivalences hold [5, Corollary 2.6]:

$$\begin{aligned} \|w\|_{H^s(\Omega)} \leq |||w||| &\lesssim \|w\|_{H^s(\Omega)}, & \text{if } s \in (0, 1/2), \\ |w|_{H^s(\Omega)} \leq |||w||| &\lesssim |w|_{H^s(\Omega)}, & \text{if } s \in (1/2, 1). \end{aligned}$$

When $s = 1/2$, since Hardy's inequality fails, it is not possible to bound the $H^{\frac{1}{2}}(\mathbb{R}^d)$ -seminorm in terms of the $H^{\frac{1}{2}}(\Omega)$ -norm for functions supported in $\bar{\Omega}$. However, for the purposes we pursue in this work, it suffices to notice that the estimate

$$|||w||| \lesssim |w|_{H^{\frac{1}{2}+\varepsilon}(\Omega)}$$

holds for all $w \in \mathbb{H}^{\frac{1}{2}+\varepsilon}(\Omega)$.

From this discussion, it follows that the energy norm may be bounded in terms of fractional-order norms on Ω . Thus, in order to estimate errors in the energy norm, we may bound errors within Ω .

3.1. Regularity. We now review some results regarding Sobolev regularity of solutions to problem (3.2) that are useful to deduce convergence rates of the finite element scheme proposed below. Regularity results for the fractional Laplacian have been recently obtained by Grubb [66] in terms of Hörmander μ -spaces [68].

The work [5] has reinterpreted these in terms of standard Sobolev spaces. The following result, see [28, 66], holds for domains with smooth boundaries, a condition that is too restrictive for a finite element analysis.

Theorem 3.1 (smooth domains). *Let $s \in (0, 1)$, Ω be a domain with $\partial\Omega \in C^\infty$, $f \in H^r(\Omega)$ for some $r \geq -s$, u be the solution of (1.4) and $\alpha = \min\{s+r, 1/2-\varepsilon\}$, with $\varepsilon > 0$ arbitrarily small. Then, $u \in \mathbb{H}^{s+\alpha}(\Omega)$ and the following regularity estimate holds:*

$$\|u\|_{\mathbb{H}^{s+\alpha}(\Omega)} \lesssim \|f\|_{H^r(\Omega)},$$

where the hidden constant depends on the domain Ω , the dimension d , s and α .

As a consequence of the previous result, we see that smoothness of the right hand side f does not ensure that solutions are any smoother than $\mathbb{H}^{s+\frac{1}{2}-\varepsilon}(\Omega)$; see also [123]. We illustrate this phenomenon with the following example.

Example 3.2 (limited regularity). We follow [62, 108] and consider $\Omega = B(0, 1) \subset \mathbb{R}^d$ and $f \equiv 1$. Then the solution to (3.2) is given by

$$(3.3) \quad u(x) = \frac{\Gamma(\frac{d}{2})}{2^{2s}\Gamma(\frac{d+2s}{2})\Gamma(1+s)}(1-|x|^2)_+^s,$$

where $t_+ = \max\{t, 0\}$.

The lack of a lifting property for the solution to (3.2) can also be explained by the fact that the eigenfunctions of this operator have reduced regularity [28, 67, 110]. This is in stark contrast with the spectral fractional Laplacian (1.6), discussed in Section 2.2, whose eigenfunctions coincide with those of the Laplacian and thus are smooth functions if the boundary of the domain is regular enough.

On the other hand, Hölder regularity results for (3.2) have been obtained in [109]. They give rise to Sobolev estimates for solutions in terms of Hölder norms of the data, that are valid for rough domains. More precisely, we have the following result; see [5, Propositions 3.6 and 3.11].

Theorem 3.3 (Lipschitz domains). *Let $s \in (0, 1)$ and Ω be a Lipschitz domain satisfying the exterior ball condition. If $s \in (0, 1/2)$, let $f \in C^{\frac{1}{2}-s}(\overline{\Omega})$; if $s = 1/2$, let $f \in L^\infty(\Omega)$; and if $s \in (1/2, 1)$, let $f \in C^\beta(\overline{\Omega})$ for some $\beta > 0$. Then, for every $\varepsilon > 0$, the solution u of (3.2) belongs to $\mathbb{H}^{s+\frac{1}{2}-\varepsilon}(\Omega)$, with*

$$\|u\|_{\mathbb{H}^{s+\frac{1}{2}-\varepsilon}(\Omega)} \lesssim \frac{1}{\varepsilon} \|f\|_\star,$$

where $\|\cdot\|_\star$ denotes the $C^{\frac{1}{2}-s}(\overline{\Omega})$, $L^\infty(\Omega)$ or $C^\beta(\overline{\Omega})$, correspondingly to whether s is smaller, equal or greater than $1/2$, and the hidden constant depends on the domain Ω , the dimension d and s .

In case $s > 1/2$, the theorem above ensures that the solution u belongs at least to $H_0^1(\Omega)$. It turns out that to prove the full $H^{s+\frac{1}{2}-\varepsilon}$ -regularity, an intermediate step is to ensure that the gradient of u is actually an L^2 -function. Following [29], this fact can be proved studying the behavior of the fractional seminorms $|\cdot|_{H^{1-s}(\Omega)}$, which usually blow up as $\delta \rightarrow 0$:

$$(3.4) \quad \lim_{\delta \rightarrow 0} \delta |w|_{H^{1-s}(\Omega)}^2 = C(d) |w|_{H^1(\Omega)}^2 \quad \forall w \in H^1(\Omega).$$

Therefore, the technique used in [5] to prove Theorem 3.3 consists of first proving that the left-hand side of (3.4) remains bounded as $\delta \rightarrow 0$ for the solution u of (3.2), whence $u \in H^1(\Omega)$, and next analyzing the regularity of the gradient of u .

As already stated in (1.7), the solution to (3.2) behaves like $\text{dist}(x, \partial\Omega)^s$ for points x close to the boundary $\partial\Omega$. This can clearly be seen in Example 3.2 and explains the reduced regularity obtained in Theorems 3.1 and 3.3. To capture this

behavior, we develop estimates in fractional weighted norms, where the weight is a power of the distance to the boundary. Following [5] we introduce the notation

$$\delta(x, x') = \min \{ \text{dist}(x, \partial\Omega), \text{dist}(x', \partial\Omega) \},$$

and, for $\ell = k + s$, with $k \in \mathbb{N}$ and $s \in (0, 1)$, and $\kappa \geq 0$, we define the norm

$$\|w\|_{H_\kappa^\ell(\Omega)}^2 := \|w\|_{H^k(\Omega)}^2 + \sum_{|\beta|=k} \iint_{\Omega \times \Omega} \frac{|D^\beta w(x) - D^\beta w(x')|^2}{|x - x'|^{d+2s}} \delta(x, x')^{2\kappa} \, dx' \, dx.$$

and the associated space

$$(3.5) \quad H_\kappa^\ell(\Omega) := \{w \in H^\ell(\Omega) : \|w\|_{H_\kappa^\ell(\Omega)} < \infty\}.$$

Although we are interested in the case $\kappa \geq 0$, we recall that in the definition of weighted Sobolev spaces $H_\kappa^k(\Omega)$, with k being a nonnegative integer, arbitrary powers of δ can be considered [77, Theorem 3.6]. On the other hand, global versions $H_\kappa^\ell(\mathbb{R}^d)$ are defined integrating in the space \mathbb{R}^d and taking δ as before, but some restrictions must be taken into account to ensure their completeness. A sufficient condition is that the weight belongs to the Muckenhoupt class $A_2(\mathbb{R}^d)$ [75]. In this context, this implies that if $|\kappa| < 1/2$ then the spaces $H_\kappa^\ell(\mathbb{R}^d)$ are complete.

Remark 3.4 (explicit solutions). In radial domains, spaces like (3.5) have been used to characterize the mapping properties of the fractional Laplacian. In particular, when Ω is the unit ball, upon defining the weight $\omega(x) = 1 - |x|^2$, an explicit eigendecomposition of the operator $w \mapsto (-\Delta)^s(\omega^s w)$ is obtained in [6, 57]. The eigenfunctions are products of solid harmonic polynomials and radial Jacobi polynomials or, in one dimension, Gegenbauer polynomials. Mapping properties of the fractional Laplacian can thus be characterized in terms of weighted Sobolev spaces defined by means of expansions on these polynomials.

The regularity in the weighted Sobolev spaces (3.5) reads as follows [5, Proposition 3.12].

Theorem 3.5 (weighted Sobolev estimate). *Let Ω be a bounded, Lipschitz domain satisfying the exterior ball condition, $s \in (1/2, 1)$, $f \in C^{1-s}(\overline{\Omega})$ and u be the solution of (3.2). Then, for every $\varepsilon > 0$ we have $u \in H_{1/2-\varepsilon}^{1+s-2\varepsilon}(\Omega)$ and*

$$\|u\|_{H_{1/2-\varepsilon}^{1+s-2\varepsilon}(\Omega)} \lesssim \frac{1}{\varepsilon} \|f\|_{C^{1-s}(\overline{\Omega})},$$

where the hidden constant depends on the domain Ω , the dimension d and s .

It is not straightforward to extend Theorem 3.5 to $s \in (0, 1/2]$ since, in this case, we cannot invoke Theorem 3.3 to obtain that the solution u belongs to $H^1(\Omega)$. Circumventing this would require to introduce a weight to obtain, for some $\kappa > 0$, that $u \in H_\kappa^{1-\varepsilon}(\Omega)$ for some $\kappa > 0$. However, for this is necessary to obtain a weighted version of (3.4) which, to the best of our knowledge, is not available in the literature. In spite of this, the numerical experiments we have carried out using graded meshes and $s \in (0, 1/2]$ show the same order of convergence as for $s \in (1/2, 1)$; see Table 1 below.

The proof of Theorem 3.5 presented in [5] also shows that, if $f \in C^\beta(\overline{\Omega})$ for some $\beta \in (0, 2 - 2s)$, $\ell < \min\{\beta + 2s, \kappa + s + 1/2\}$, then we have

$$(3.6) \quad |u|_{H_\kappa^\ell(\Omega)} \lesssim \frac{1}{(\beta + \ell - 2s)(1 + 2(\kappa - s - \ell))} |f|_{C^\beta(\overline{\Omega})},$$

where the hidden constant depends only on Ω and the dimension d . This shows that increasing the exponent κ of the weight allows for the differentiability order ℓ to increase as well. In principle, there is no restriction on κ above; however, in the next subsection we exploit this weighted regularity by introducing approximations

on a family of graded meshes. There we show that the order of convergence (with respect to the number of degrees of freedom) is only incremented as long as $\kappa < 1/2$.

3.2. FEM: A Priori Error Analysis. The numerical approximation of the solution to (3.2) presents an immediate difficulty: the kernel of the bilinear form (3.1) is singular, and consequently, special care must be taken. For this reason, most existing approaches restrict themselves to the one dimensional case ($d = 1$). Explorations in this direction can be found using finite elements [51], finite differences [69] and Nyström methods [6]. For several dimensions the literature is rather scarce. A Monte Carlo algorithm that avoids dealing with the singular kernel was proposed in [79]; we also refer the reader to the recent work [8], that builds on the results from [4, 5] to explore the connection between the integral fractional Laplacian with boundary integral operators to develop efficient finite element techniques.

Here we present a direct finite element approximation in arbitrary dimensions. Following §2.4 we denote by \mathcal{T} a conforming and shape regular mesh of Ω , consisting of simplicial elements K of diameter bounded by h . In order to present a unified approach for the whole range $s \in (0, 1)$ we only consider approximations of (3.2) by continuous functions. For $s < 1/2$ it is possible to consider piecewise constants, but we do not explore this here. With the finite element space $\mathbb{U}(\mathcal{T})$ defined as in (2.10), the finite element approximation of (3.2) is then the unique solution to the problem: find $U \in \mathbb{U}(\mathcal{T})$ such that

$$(3.7) \quad \llbracket U, W \rrbracket = \langle f, W \rangle \quad \forall W \in \mathbb{U}(\mathcal{T}).$$

From this formulation it immediately follows that U is the projection (in the energy norm) of u onto $\mathbb{U}(\mathcal{T})$. Consequently, we have a Céa-like best approximation result

$$\| \| u - U \| \| = \inf_{W \in \mathbb{U}(\mathcal{T})} \| \| u - W \| \| .$$

Thus, in order to obtain a priori rates of convergence, it just remains to bound the energy-norm distance between the discrete spaces and the solution. One difficult aspect of dealing with fractional seminorms is that they are not additive with respect to domain decompositions. Nevertheless, it is possible to localize these norms [61]: for all $w \in H^s(\Omega)$ we have

$$|w|_{H^s(\Omega)}^2 \leq \sum_{K \in \mathcal{T}} \left[\int_K \int_{S_K} \frac{|w(x) - w(x')|^2}{|x - x'|^{d+2s}} dx' dx + \frac{C(d, \sigma)}{sh_K^{2s}} \|w\|_{L^2(K)}^2 \right],$$

where S_K is the patch associated with $K \in \mathcal{T}$ and σ denotes the shape-regularity parameter of the mesh \mathcal{T} . From this inequality it follows that, to obtain a priori error estimates, it suffices to compute interpolation errors over the set of patches $\{K \times S_K\}_{K \in \mathcal{T}}$. The reduced regularity of solutions implies that we need to resort to quasi-interpolation operators; we work with the Scott-Zhang operator $\Pi_{\mathcal{T}}$ [114]. Local stability and approximation properties of this operator were studied by Ciarlet Jr. in [46].

Proposition 3.6 (quasi-interpolation estimate). *Let $K \in \mathcal{T}$, $\max\{1/2, s\} < \ell \leq 2$, $s \in (0, 1)$, and $\Pi_{\mathcal{T}}$ be the Scott-Zhang operator. If $w \in H^\ell(\Omega)$, then*

$$\int_K \int_{S_K} \frac{|(w - \Pi_{\mathcal{T}}w)(x) - (w - \Pi_{\mathcal{T}}w)(x')|^2}{|x - x'|^{d+2s}} dx' dx \lesssim h_K^{2\ell-2s} |w|_{H^\ell(S_K)}^2,$$

where the hidden constant depends on d, σ, ℓ and blows up as $s \uparrow 1$.

The interpolation estimate of Proposition 3.6 shows that, if the meshsize is sufficiently small and we set $\varepsilon \approx |\log h|^{-1}$ in the estimates of Theorems 3.3 and 3.5; see [5, Theorem 4.7] we can deduce an a priori error bound in the energy norm .

Theorem 3.7 (energy error estimate for quasi-uniform meshes). *Let u denote the solution to (3.2) and denote by $U \in \mathbb{U}(\mathcal{T})$ the solution of the discrete problem (3.7), computed over a mesh \mathcal{T} consisting of elements with maximum diameter h . Under the hypotheses of Theorem 3.3 we have*

$$\|u - U\| \lesssim h^{\frac{1}{2}} |\log h| \|f\|_{\star},$$

where the hidden constant depends on Ω , s and σ , and $\|\cdot\|_{\star}$ denotes the $C^{\frac{1}{2}-s}(\overline{\Omega})$, $L^{\infty}(\Omega)$ or $C^{\beta}(\overline{\Omega})$, correspondingly to whether s is smaller, equal or greater than $1/2$.

This estimate hinges on the regularity of solutions provided by Theorem 3.3, and thus it depends on Hölder bounds for the data. We now turn our attention to obtaining a priori error estimates in the $L^2(\Omega)$ -norm. Using Theorem 3.1, an Aubin-Nitsche duality argument can be carried out. The proof of the following proposition follows the steps outlined in [28, Proposition 4.3].

Proposition 3.8 (L^2 -error estimate). *Let u denote the solution to (3.2) and denote by $U \in \mathbb{U}(\mathcal{T})$ the solution of the discrete problem (3.7), computed over a mesh \mathcal{T} consisting of elements with maximum diameter h . Under the hypotheses of Theorem 3.1 we have*

$$\|u - U\|_{L^2(\Omega)} \lesssim h^{\alpha+\beta} \|f\|_{H^r(\Omega)},$$

where $\alpha = \min\{s + r, 1/2 - \varepsilon\}$, $\beta = \min\{s, 1/2 - \varepsilon\}$, $\varepsilon > 0$ may be taken arbitrarily small and the hidden constant depends on Ω , s , d , σ , α and blows up when $\varepsilon \rightarrow 0$.

Finally, for $s \in (1/2, 1)$ and $d = 2$, we take advantage of Theorem 3.5, from which further information about the boundary behavior of solutions is available. We propose a standard procedure often utilized in connection with corner singularities or boundary layers arising in convection-dominated problems. An increased rate of convergence is achieved by resorting to a priori adapted meshes. To obtain interpolation estimates in the weighted fractional Sobolev spaces defined by (3.5), we introduce the following Poincaré inequality [5, Proposition 4.8].

Proposition 3.9 (weighted fractional Poincaré inequality). *Let $s \in (0, 1)$, $\kappa \in [0, s)$ and a domain S which is star-shaped with respect to a ball. Then, for every $w \in H_{\kappa}^s(S)$, it holds*

$$\|w - \bar{w}\|_{L^2(S)} \lesssim d_S^{s-\kappa} |w|_{H_{\kappa}^s(S)},$$

where $\bar{w} = \int_S w \, dx$, $d_S = \text{diam}(S)$ and the hidden constant depends on the chunkiness parameter of S and the dimension d .

This inequality yields sharp quasi-interpolation estimates near the boundary of the domain, where the weight δ involved in (3.5) degenerates. Such bounds in turn lead to estimates for the Scott-Zhang operator in weighted fractional spaces. We exploit them for two-dimensional problems ($d = 2$) by constructing *graded* meshes as in [65, Section 8.4]. In addition to shape regularity, we assume that our meshes \mathcal{T} satisfy the following property: there is a number $\mu \geq 1$ such that given a meshsize parameter h and $K \in \mathcal{T}$, we have

$$(3.8) \quad h_K \leq C(\sigma) \begin{cases} h^{\mu}, & K \cap \partial\Omega \neq \emptyset, \\ h \text{dist}(K, \partial\Omega)^{(\mu-1)/\mu}, & K \cap \partial\Omega = \emptyset. \end{cases}$$

where the constant $C(\sigma)$ depends only on the shape regularity constant σ of the mesh \mathcal{T} . The parameter μ relates the meshsize h to the number of degrees of

freedom because (recall that $d = 2$)

$$\#\mathcal{T} \approx \begin{cases} h^{-2}, & \mu < 2, \\ h^{-2}|\log h|, & \mu = 2, \\ h^{-\mu}, & \mu > 2. \end{cases}$$

It is now necessary to relate the parameter μ with the exponent κ of the weight δ in estimate (3.6). Increasing the parameter μ corresponds to raising κ and thereby allowing an increase of the differentiability order ℓ . However, if $\mu > 2$ this gain is compensated by a growth in the number of degrees of freedom. Following [5], it turns out that the optimal parameter is $\mu = 2$ and we have the following result.

Theorem 3.10 (energy error estimates for graded meshes). *Let $\Omega \subset \mathbb{R}^2$ and $U \in \mathbb{U}(\mathcal{T})$ be the solution to (3.7), computed over a mesh \mathcal{T} that satisfies (3.8) with $\mu = 2$. In the setting of Theorem 3.5 we have*

$$\|u - U\| \lesssim (\#\mathcal{T})^{-\frac{1}{2}} |\log(\#\mathcal{T})| \|f\|_{C^{1-s}(\bar{\Omega})},$$

where the hidden constant depends on σ and blows up as $s \rightarrow 1/2$.

3.3. Implementation. Let us now discuss key details about the finite element implementation of (3.2) for $d = 2$. If $\{\phi_v\}$ are the nodal piecewise linear basis functions of $\mathbb{U}(\mathcal{T})$, defined as in (2.10), then the entries of the stiffness matrix $\mathcal{A} = (\mathcal{A}_{vw})$ are

$$\mathcal{A}_{vw} = \llbracket \phi_v, \phi_w \rrbracket = \frac{C(d, s)}{2} \iint_Q \frac{(\phi_v(x) - \phi_v(x'))(\phi_w(x) - \phi_w(x'))}{|x - x'|^{2+2s}} dx' dx.$$

Two numerical difficulties — coping with integration on unbounded domains and handling the *non-integrable* singularity of the kernel — seem to discourage a direct finite element approach. However, borrowing techniques from the boundary element method [113], it is possible to compute accurately the entries of the matrix \mathcal{A} . We next briefly outline the main steps of this procedure. For full details, we refer to [4] where a finite element code to solve (3.2) is documented.

The integrals involved in the computation of \mathcal{A} should be carried over \mathbb{R}^2 . For this reason it is convenient to consider a ball B containing Ω and such that the distance from $\bar{\Omega}$ to B^c is an arbitrary positive number. This is needed in order to avoid difficulties caused by lack of symmetry when dealing with the integral over Ω^c when Ω is not a ball. Together with B , we introduce an auxiliary triangulation \mathcal{T}_A on $B \setminus \Omega$ such that the complete triangulation \mathcal{T}_B over B (that is $\mathcal{T}_B = \mathcal{T} \cup \mathcal{T}_A$) remains admissible and shape-regular.

We define, for $1 \leq \ell, m \leq \#\mathcal{T}_B$ and $K_\ell, K_m \in \mathcal{T}_B$,

$$(3.9) \quad \begin{aligned} I_{\ell, m}^{v, w} &:= \int_{K_\ell} \int_{K_m} \frac{(\phi_v(x) - \phi_v(x'))(\phi_w(x) - \phi_w(x'))}{|x - x'|^{2+2s}} dx' dx, \\ J_\ell^{v, w} &:= \int_{K_\ell} \int_{B^c} \frac{\phi_v(x)\phi_w(x)}{|x - x'|^{2+2s}} dx' dx, \end{aligned}$$

whence we may write

$$\mathcal{A}_{vw} = \frac{C(d, s)}{2} \sum_{\ell=1}^{\#\mathcal{T}_B} \left(\sum_{m=1}^{\#\mathcal{T}_B} I_{\ell, m}^{v, w} + 2J_\ell^{v, w} \right).$$

We reiterate that computing the integrals $I_{\ell, m}^{v, w}$ and $J_\ell^{v, w}$ is challenging for different reasons: the former involves a singular integrand if $\bar{K}_\ell \cap \bar{K}_m \neq \emptyset$, while the latter needs to be calculated in an unbounded domain.

We first tackle the computation of $I_{\ell, m}^{v, w}$ in (3.9). If \bar{K}_ℓ and \bar{K}_m do not touch, then the integrand is a regular function and can be integrated numerically in a

standard fashion. On the other hand, if $\overline{K_\ell} \cap \overline{K_m} \neq \emptyset$, then $I_{\ell,m}^{v,w}$ bears some resemblances to typical integrals appearing in the boundary element method. Indeed, the quadrature rules we employ are analogous to the ones presented in [113, Chapter 5]. Basically, the scheme consists of the following steps:

- Consider parametrizations $\chi_\ell: \hat{K} \rightarrow K_\ell$ and $\chi_m: \hat{K} \rightarrow K_m$ such that the edge or vertex shared by K_ℓ and K_m is the image of the same edge/vertex in the reference element \hat{K} . If K_ℓ and K_m coincide, simply use the same parametrization twice.
- Decompose the integration domain into certain subsimplices and then utilize Duffy-type transformations to map these subdomains into the four-dimensional unit hypercube.
- Since the Jacobian of these Duffy transformations is regularizing, each of the integrals may be separated into two parts: a highly singular but explicitly integrable part and a smooth, numerically tractable part.

The second difficulty lies in the calculation of $J_\ell^{v,w}$, namely, dealing with the unbounded domain B^c . We write

$$J_\ell^{v,w} = \int_{K_\ell} \phi_v(x) \phi_w(x) \varrho(x) \, dx, \quad \varrho(x) := \int_{B^c} \frac{1}{|x - x'|^{2+2s}} \, dx',$$

and realize that we need to accurately compute $\varrho(x)$ at each quadrature point $x \in K_\ell \cap \bar{\Omega}$. To do so, there are two properties we can take advantage of: the radially of ϱ and the fact that ϱ is smooth up to the boundary of Ω because, for $x \in \bar{\Omega}$ and $x' \in B^c$ $|x - x'| > \text{dist}(\bar{\Omega}, B^c) > 0$. Therefore, the values of ϱ at quadrature nodes can be precomputed with an arbitrary degree of precision.

3.4. Numerical Experiments. We present the outcome of two experiments posed in $\Omega = B(0, 1) \subset \mathbb{R}^2$.

3.4.1. Rate of Convergence in Energy Norm. Following Example 3.2 we have that, if $f \equiv 1$, then the solution to (3.2) is given by (3.3). Table 1 shows computational rates of convergence in the energy norm for several values of s , both for uniform and graded meshes. These rates are in agreement with those predicted by Theorems 3.7 and 3.10. Moreover, we observe an increased order of convergence for $s \leq 1/2$ and graded meshes which is not accounted for in Theorem 3.10.

Value of s	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
Uniform meshes	0.497	0.496	0.498	0.500	0.501	0.505	0.504	0.503	0.532
Graded meshes	1.066	1.040	1.019	1.002	1.066	1.051	0.990	0.985	0.977

TABLE 1. Example 3.4.1: Computational rates of convergence for (3.2) posed in the unit ball with right-hand side $f \equiv 1$. Errors are measured in the energy norm with respect to the meshsize parameter h . The second row corresponds to uniform meshes, while the third to graded meshes, with $\mu = 2$ in (3.8).

3.4.2. Rate of Convergence in L^2 -Norm. Remark 3.4 states that a family of explicit solutions for (3.2) is available in the unit ball. A subclass of solutions in that family may be expressed in terms of the Jacobi polynomials $P_k^{(\alpha,\beta)}: [-1, 1] \rightarrow \mathbb{R}$. We set

$$f(x) = \left(\frac{\Gamma(3+s)}{2^{1-s}} \right)^2 P_2^{(s,0)}(2|x|^2 - 1),$$

so that the solution to (3.2) is given by [57, Theorem 3]

$$u(x) = (1 - |x|^2)_+^s P_2^{(s,0)}(2|x|^2 - 1).$$

We compute the orders of convergence in $L^2(\Omega)$ for $s \in \{0.25, 0.75\}$; according to Proposition 3.8, it is expected to have order of convergence 0.75 for $s = 0.25$ and 1 for $s = 0.75$ with respect to the meshsize h . The results, summarized in Table 2, agree with the predicted rates of convergence.

h	$s = 0.25$	$s = 0.75$
0.0383	0.0801	0.01740
0.0331	0.0698	0.01388
0.0267	0.0605	0.01104
0.0239	0.0556	0.00965
0.0218	0.0513	0.00849

TABLE 2. Example 3.4.2: Errors in the L^2 -norm for $s = 0.25$ and $s = 0.75$. The estimated orders of convergence with respect to the meshsize are, respectively, 0.7669 and 1.2337.

3.5. FEM: A Posteriori Error Analysis. Since solutions of (3.2) have reduced regularity, and assembling the stiffness matrix \mathcal{A} entails a rather high computational cost, it is of interest to devise suitable AFEMs. We now present a posteriori error estimates of residual type and ensuing AFEM; we follow [101].

We stress that, in contrast to the case $s = 1$, $(-\Delta)^s U$ does not have jumps on inter-element boundaries and does not vanish within elements for piecewise linear U . We estimate the energy error $\|u - U\|$ in terms of the residual $\mathcal{R} := f - (-\Delta)^s U$ in $\mathbb{H}^{-s}(\Omega)$. To do so, we need to address two important issues: localization of the norm in $\mathbb{H}^{-s}(\Omega)$ and a practical computation of \mathcal{R} .

To localize fractional norms we deviate from [61] and perform a decomposition on stars $S_v = \text{supp } \phi_v$, the support of the basis functions ϕ_v associated with node v and diameter h_v . Exploiting the partition of unity property $\sum_v \phi_v = 1$, and Galerkin orthogonality $\llbracket u - U, \phi_v \rrbracket = 0$ for all $v \in \Omega$, we can write, for every $w \in \mathbb{H}^s(\Omega)$

$$\llbracket u - U, w \rrbracket = \langle \mathcal{R}, w \rangle = \sum_v \langle \mathcal{R}, w \phi_v \rangle = \sum_v \langle \mathcal{R}, (w - \bar{w}_v) \phi_v \rangle = \sum_v \langle (\mathcal{R} - \bar{\mathcal{R}}_v) \phi_v, w - \bar{w}_v \rangle,$$

where $\bar{w}_v \in \mathbb{R}$ are weighted mean values computed as $\bar{w}_v = 0$ provided $v \in \partial\Omega$ and, otherwise,

$$\bar{w}_v := \frac{\langle w, \phi_v \rangle}{\langle \phi_v, 1 \rangle} \quad \forall v \in \Omega.$$

The values of $\bar{\mathcal{R}}_v \in \mathbb{R}$ are yet to be chosen. We see that $\mathcal{R} = \sum_v (\mathcal{R} - \bar{\mathcal{R}}_v) \phi_v$ where each term $(\mathcal{R} - \bar{\mathcal{R}}_v) \phi_v \in \mathbb{H}^{-s}(S_v)$ has support in S_v . We have the following two estimates for dual norms proved in [101, Lemmas 1 and 2].

Lemma 3.11 (localized upper bound of dual norm). *Let $\mathcal{G} \in \mathbb{H}^{-s}(\Omega)$ be decomposed as $\mathcal{G} = \sum_v g_v$ with $g_v \in \mathbb{H}^{-s}(S_v)$ vanishing outside S_v . We then have for $0 < s < 1$*

$$\|\mathcal{G}\|_{\mathbb{H}^{-s}(\Omega)}^2 \leq (d+1) \sum_v \|g_v\|_{\mathbb{H}^{-s}(S_v)}^2.$$

A key practical issue is then how to evaluate $\|g_v\|_{\mathbb{H}^{-s}(S_v)}$ for $g_v = (\mathcal{R} - \bar{\mathcal{R}}_v) \phi_v$. For second order operators, \mathcal{R} splits into an L^2 -component in element interiors (provided $f \in L^2(\Omega)$) and a singular component supported on element boundaries. In contrast, the residual $\mathcal{R} = f - (-\Delta)^s U$ does not have a singular component and its absolutely continuous part is not always in $L^2(\Omega)$ for $s \in [3/4, 1)$ no matter how smooth f might be. This is related to singularities of $(-\Delta)^s \phi_v(x)$ as x tends to the

skeleton of $S_{\mathbf{v}}$ because $\phi_{\mathbf{v}}$ is continuous, piecewise linear. Given $t \geq 0$ and $p > 1$, let

$$\widetilde{W}_p^t(\Omega) := \{w|_{\Omega} : w \in W_p^t(\mathbb{R}^d), w|_{\mathbb{R}^d \setminus \Omega} = 0\}.$$

Using that [121, Theorem XI.2.5]

$$(-\Delta)^s : \widetilde{W}_p^t(\Omega) \rightarrow W_p^{t-2s}(\Omega),$$

is a continuous pseudo-differential operator of order $2s$, and $\phi_{\mathbf{v}} \in \widetilde{W}_p^{1+\frac{1}{p}-\varepsilon}(\Omega)$ for any $\varepsilon > 0$, we deduce

$$(-\Delta)^s U \in L^p(\Omega) \quad \frac{1}{p} > 2s - 1.$$

This motivates the following estimate, whose proof is given in [101, Lemma 2].

Lemma 3.12 (upper bound of local dual norm). *Let $g_{\mathbf{v}} \in L^p(S_{\mathbf{v}})$ satisfy $\int_{S_{\mathbf{v}}} g_{\mathbf{v}} \, dx = 0$ for each $\mathbf{v} \in \Omega$. If $0 < s < 1$ and $1 \leq p < \infty$ satisfies $\frac{1}{p} < \frac{s}{d} + \frac{1}{2}$, then*

$$\|g_{\mathbf{v}}\|_{\mathbb{H}^{-s}(S_{\mathbf{v}})} \lesssim h_{\mathbf{v}}^{s+\frac{d}{2}-\frac{d}{p}} \|g_{\mathbf{v}}\|_{L^p(S_{\mathbf{v}})}.$$

However, to be able to apply Lemma 3.12 the Lebesgue exponent p must satisfy

$$2s - 1 < \frac{1}{p} < \frac{s}{d} + \frac{1}{2}.$$

We note that for $s < \frac{3}{4}$ we can choose $p = 2$ for any dimension d . However, for $\frac{3}{4} \leq s < 1$ we need to take $1 < p < 2$. For $d = 1, 2$ this condition is satisfied for any $s < 1$, but for $d = 3$ we have the unfortunate constraint $s < \frac{9}{10}$.

We can now choose $\bar{\mathcal{R}}_{\mathbf{v}}$ as

$$\bar{\mathcal{R}}_{\mathbf{v}} := \frac{\int_{S_{\mathbf{v}}} \mathcal{R} \phi_{\mathbf{v}} \, dx}{\int_{S_{\mathbf{v}}} \phi_{\mathbf{v}} \, dx}, \quad \mathbf{v} \in \Omega$$

and $\bar{\mathcal{R}}_{\mathbf{v}} = 0$ otherwise, so that the local contributions satisfy $\int_{S_{\mathbf{v}}} (\mathcal{R} - \bar{\mathcal{R}}_{\mathbf{v}}) \phi_{\mathbf{v}} \, dx = 0$ and we can then apply the bound of Lemma 3.12. The following upper a posteriori error estimate is derived in [101, Theorem 1].

Theorem 3.13 (upper a posteriori bound). *Let $f \in L^p(\Omega)$ and $1 < p < \infty, 0 < s < 1$ satisfy the restriction $2s - 1 < \frac{1}{p} < \frac{s}{d} + \frac{1}{2}$, then*

$$\|u - U\|_{\mathbb{H}^{-s}(\Omega)}^2 \lesssim \sum_{\mathbf{v}} h_{\mathbf{v}}^{2(s+\frac{d}{2}-\frac{d}{p})} \|(\mathcal{R} - \bar{\mathcal{R}}_{\mathbf{v}}) \phi_{\mathbf{v}}\|_{L^p(S_{\mathbf{v}})}^2.$$

This error analysis has two pitfalls. The first one, alluded to earlier, is a restriction on s for $d > 2$. The second one is the actual computation of $(-\Delta)^s \phi_{\mathbf{v}}(x)$ for $d > 1$, which is problematic due to its singular behavior as x tends to the skeleton on \mathcal{T} . This is doable for $d = 1$ and we refer to [101, Section 8] for details and numerical experiments. This topic is obviously open for improvement.

3.6. Extensions and Applications. We conclude the discussion by mentioning extensions and applications of this approach:

- *Eigenvalue problems:* The eigenvalue problem for the integral fractional Laplacian arises, for example, in the study of fractional quantum mechanics [81]. As already mentioned, a major difference between the spectral fractional Laplacian and the integral one is that, for the second one, eigenfunctions have reduced regularity. In [28], conforming finite element approximations were analyzed and it was shown that the Babuška-Osborn theory [14] holds in this context. Regularity results for the eigenfunctions are derived under the assumption that the domain is Lipschitz and satisfies the exterior ball condition. Numerical evidence on the

non-convex domain $\Omega = (-1, 1)^2 \setminus [0, 1]^2$ indicates that the first eigenfunction is as regular as the first one on any smooth domain. This is in contrast with the Laplacian.

- *Time dependent problems:* In [3], problem (2.17) with $\gamma \in (0, 2]$ and the integral definition of $(-\Delta)^s$ was considered. Regularity of solutions was studied and a discrete scheme was proposed and analyzed. The method is based on a standard Galerkin finite element approximation in space, as described here, while in time a convolution quadrature approach was used [70, 82].
- *Non-homogeneous Dirichlet conditions:* An interpretation of a non-homogeneous Dirichlet condition g for the integral fractional Laplacian is given by using (1.3) upon extension by g over Ω^c . In [7] a mixed method for this problem is proposed; it is based on the weak enforcement of the Dirichlet condition and the incorporation of a certain non-local normal derivative as a Lagrange multiplier. This non-local derivative is interpreted as a non-local flux between Ω and Ω^c [53].
- *Non-local models for interface problems:* Consider two materials with permittivities/diffusivities of opposite sign, and separated by an interface with a corner. Strong singularities may appear in the classical (local) models derived from electromagnetics theory. In fact, the problem under consideration is of Fredholm type if and only if the quotient between the value of permittivities/diffusivities taken from both sides of the interface lies outside a so-called critical interval, which always contains the value -1 . In [27] a non-local interaction model for the materials is proposed. Numerical evidence indicates that the non-local model may reduce the critical interval and that solutions are more stable than for the local problem.

4. DUNFORD-TAYLOR APPROACH FOR SPECTRAL AND INTEGRAL LAPLACIANS

In this section we present an alternative approach to the ones developed in the previous sections. It relies on the Dunford-Taylor representation (1.11)

$$u = (-\Delta)^{-s} f = \frac{\sin(s\pi)}{\pi} \int_0^\infty \mu^{-s} (\mu - \Delta)^{-1} f \, d\mu$$

for the spectral fractional Laplacian (1.6). For the integral fractional Laplacian (1.2), instead, it hinges on the equivalent representation (1.13) of (1.12):

$$(4.1) \quad \int_{\mathbb{R}^d} |\xi|^s \mathcal{F}(\tilde{u}) |\xi|^s \overline{\mathcal{F}(\tilde{w})} \, d\xi = \frac{2 \sin(s\pi)}{\pi} \int_0^\infty \mu^{1-2s} \int_{\Omega} ((-\Delta)(I - \mu^2 \Delta)^{-1} \tilde{u}(x)) w(x) \, dx \, d\mu.$$

In (4.1), the operators $-\Delta$ and $I - \mu^2 \Delta$ are defined over \mathbb{R}^d , something to be made precise in Theorem 4.5.

In each case, the proposed method is proved to be efficient on *general Lipschitz domains* $\Omega \subset \mathbb{R}^d$. They rely on sinc quadratures and on finite element approximations of the resulting integrands at each quadrature points. While (1.11) allows for a direct approximation of the solution, the approximation of (1.13) leads to a non-conforming method where the action of the stiffness matrix on a vector is approximated.

We recall that the functional spaces $\mathbb{H}^s(\Omega)$ are defined in (1.5) for $s \in [0, 3/2)$. We now extend the definition $\mathbb{H}^s(\Omega) = H^s(\Omega) \cap H_0^1(\Omega)$ for $s \in (1, 2]$.

4.1. Spectral Laplacian. We follow [25, 26] and describe a method based on the Balakrishnan representation (1.11). In order to simplify the notation, we set $L := -\Delta : \mathcal{D}(L) \rightarrow L^2(\Omega)$ and define the domain of L^r , for $r \in \mathbb{R}$, to be

$$\mathcal{D}(L^r) := \{v \in L^2(\Omega) : L^r v \in L^2(\Omega)\};$$

this is a Banach space equipped with the norm

$$\|v\|_{\mathcal{D}(L^r)} := \|L^r v\|_{L^2(\Omega)}.$$

We also define the solution operator $T : H^{-1}(\Omega) \rightarrow H_0^1(\Omega)$ by $TF := v$, where for $F \in H^{-1}(\Omega)$, $v \in H_0^1(\Omega)$ is the unique solution of

$$\int_{\Omega} \nabla v \cdot \nabla w \, dx = F(v), \quad \forall w \in H_0^1(\Omega).$$

This definition directly implies that $\mathcal{D}(L) = \text{range}(T|_{L^2(\Omega)})$.

4.1.1. Finite Element Discretization. For simplicity, we assume that the domain Ω is polytopal so that it can be partitioned into a conforming subdivision \mathcal{T} . We recall that $\mathbb{U}(\mathcal{T}) \subset H_0^1(\Omega)$ stands for the subspace of globally continuous piecewise linear polynomials with respect to \mathcal{T} ; see Section 2.4. We denote by $\Pi_{\mathcal{T}}$ the $L^2(\Omega)$ -orthogonal projection onto $\mathbb{U}(\mathcal{T})$ and by $L_{\mathcal{T}} : \mathbb{U}(\mathcal{T}) \rightarrow \mathbb{U}(\mathcal{T})$ the finite element approximation of L , i.e., for $V \in \mathbb{U}(\mathcal{T})$, $L_{\mathcal{T}}V \in \mathbb{U}(\mathcal{T})$ solves

$$\int_{\Omega} (L_{\mathcal{T}}V)W \, dx = \int_{\Omega} \nabla V \cdot \nabla W \, dx, \quad \forall W \in \mathbb{U}(\mathcal{T}).$$

We finally denote by $T_{\mathcal{T}}$ the inverse of $L_{\mathcal{T}}$, the finite element solution operator, and by h the maximum diameter of elements in \mathcal{T} .

With these notations, we are in the position to introduce the finite element approximation $U \in \mathbb{U}(\mathcal{T})$ of $u = (-\Delta)^{-s}f$ (see (1.11)):

$$(4.2) \quad U := \frac{\sin(s\pi)}{\pi} \int_0^{\infty} \mu^{-s} (\mu + L_{\mathcal{T}})^{-1} \Pi_{\mathcal{T}} f \, d\mu.$$

The efficiency of the approximation of u by U depends on the efficiency of the finite element solver $(T_{\mathcal{T}}\Pi_{\mathcal{T}})f$ (i.e. for the standard Laplacian), which is dictated by the regularity of Tf . This regularity aspect has been intensively discussed in the literature [15, 21, 50, 71, 74, 93]. In this exposition, we make the following general assumption.

Definition 4.1 (elliptic regularity). We say that T satisfies a pick-up regularity of index $0 < \alpha \leq 1$ on Ω if for $0 \leq r \leq \alpha$, the operator T is an isomorphism from $\mathbb{H}^{-1+r}(\Omega)$ to $\mathbb{H}^{1+r}(\Omega)$.

Notice that $\alpha = 1$ when Ω is convex, whence this definition extends (2.8) to general Lipschitz domains.

Assuming a pick-up regularity of index α , for any $r \in [0, 1]$, we have

$$\|Tw - T_{\mathcal{T}}\Pi_{\mathcal{T}}w\|_{\mathbb{H}^r(\Omega)} \lesssim h^{2\alpha_*} \|Tw\|_{\mathbb{H}^{\alpha+1}(\Omega)} \lesssim h^{2\alpha_*} \|w\|_{\mathbb{H}^{\alpha-1}(\Omega)}$$

where $\alpha_* := \frac{1}{2}(\alpha + \min(1-r, \alpha))$. The proof of the above estimate is classical and is based on a duality argument (Nitsche's trick); see e.g. [26, Lemma 6.1]. Notice that $\alpha_* < \alpha$ when $1 - \alpha < r$, i.e., the error is measured with regularity index too large to take full advantage of the pick-up regularity in the duality argument.

We expect that approximation (4.2) of the fractional Laplacian problem delivers the *same* rate of convergence

$$(4.3) \quad \|u - U\|_{\mathbb{H}^r(\Omega)} \lesssim h^{2\alpha_*} \|u\|_{\mathbb{H}^{1+\alpha}(\Omega)} \lesssim h^{2\alpha_*} \|f\|_{\mathbb{H}^{1+\alpha-2s}(\Omega)},$$

but the function U in (4.2) is well defined provided $f \in L^2(\Omega)$, i.e. $1 + \alpha - 2s \geq 0$.

Before describing the finite element approximation result, we make the following comments. The solution $u = L^{-s}f$ belongs to $\mathcal{D}(L^{(1+\alpha)/2})$ provided that $f \in \mathcal{D}(L^{(1+\alpha)/2-s})$. Hence, estimates such as (4.3) rely on the characterization of $\mathcal{D}(L^{r/2})$ for $r \in (0, 1 + \alpha]$. For $0 \leq r \leq 1$, the spaces $\mathcal{D}(L^{r/2})$ and $\mathbb{H}^r(\Omega)$ are equivalent, as they are both scale spaces which coincide at $r = 0$ and $r = 1$.

Furthermore, assuming an elliptic regularity pick-up of index $0 < \alpha \leq 1$, this characterization extends up to $1 + \alpha$ [26, Theorem 6.4 and Remark 4.2].

Theorem 4.2 (finite element approximation). *Assume that T satisfies a pick-up regularity of index $\alpha \in (0, 1]$ on Ω . Given $r \in [0, 1]$ with $r \leq 2s$, set $\gamma := \max\{r + 2\alpha_* - 2s, 0\}$ and $\alpha_* := \frac{1}{2}(\alpha + \min(1 - r, \alpha))$. If $f \in \mathbb{H}^\delta(\Omega)$ for $\delta \geq \gamma$, then*

$$\|u - U\|_{\mathbb{H}^r(\Omega)} \leq C_h h^{2\alpha_*} \|f\|_{\mathbb{H}^\delta(\Omega)},$$

where

$$C_h \lesssim \begin{cases} \log(2/h), & \text{when } \delta = \gamma \text{ and } r + 2\alpha_* \geq 2s, \\ 1, & \text{when } \delta > \gamma \text{ and } r + 2\alpha_* \geq 2s, \\ 1, & \text{when } \delta = 0 \text{ and } 2s > r + 2\alpha_*. \end{cases}$$

4.2. Sinc Quadrature. It remains to put in place a sinc quadrature [84] to approximate the integral in (4.2). We use the change of variable $\mu = e^y$ so that

$$U = \frac{\sin(s\pi)}{\pi} \int_{-\infty}^{\infty} e^{(1-s)y} (e^y + L_{\mathcal{T}})^{-1} \Pi_{\mathcal{T}} f \, dy.$$

Given $k > 0$, we set

$$N_+ := \left\lceil \frac{\pi^2}{4sk^2} \right\rceil, \quad N_- := \left\lceil \frac{\pi^2}{4(1-s)k^2} \right\rceil, \quad \text{and} \quad y_\ell := k\ell,$$

and define the sinc quadrature approximation of U by

$$(4.4) \quad U^k := \frac{\sin(s\pi)}{\pi} k \sum_{\ell=-N_-}^{N_+} e^{(1-s)y_\ell} (e^{y_\ell} + L_{\mathcal{T}})^{-1} \Pi_{\mathcal{T}} f.$$

The sinc quadrature consists of uniformly distributed quadrature points in the y variable. The relations $N_+, N_- \approx k^{-2}$ balance the quadrature error on $(-N_-, N_+)$ with the truncation error due to discarding the integration domain $(-\infty, -N_-) \cup (N_+, \infty)$. Moreover, the dependency in s makes the sinc quadrature more robust when s is close to 0 or 1.

The decay when $|z| \rightarrow +\infty$ and holomorphic properties of the integrand $z^{-s}(z - L)^{-1}$ in the Dunford-Taylor representation (1.10) guarantee the *exponential convergence* of the sinc quadrature [26, Theorem 7.1].

Theorem 4.3 (sinc quadrature). *For $r \in [0, 1]$, we have*

$$\|U - U^k\|_{\mathbb{H}^r(\Omega)} \lesssim e^{-\pi^2/(2k)} \|f\|_{\mathbb{H}^r(\Omega)}.$$

This estimate is not optimal in regards to the regularity required on f . In fact, the analysis provided in [26] can be easily modified to yield

$$(4.5) \quad \|U - U^k\|_{\mathbb{H}^r(\Omega)} \lesssim e^{-\pi^2/(2k)} \|f\|_{\mathbb{H}^\delta(\Omega)},$$

provided $f \in \mathbb{H}^\delta(\Omega)$ with $\delta > r - 2s$ if $r - 2s \geq 0$ and $\delta = 0$ otherwise. This will be the subject of a future note presently in preparation.

Remark 4.4 (implementation). The method based on (4.4) requires $N_+ + N_- + 1$ independent standard Laplacian finite element solves for each quadrature points y_ℓ :

$$V^\ell \in \mathbb{U}(\mathcal{T}) : \quad e^{y_\ell} \int_{\Omega} V^\ell W \, dx + \int_{\Omega} \nabla V^\ell \cdot \nabla W \, dx = \int_{\Omega} f W \, dx \quad \forall W \in \mathbb{U}(\mathcal{T}),$$

which are then aggregated to yield U^k :

$$U^k = \frac{\sin(s\pi)}{\pi} k \sum_{\ell=-N_-}^{N_+} e^{(1-s)y_\ell} V^\ell.$$

Implementation of this algorithm starting from a finite element solver for the Poisson problem is straightforward. Numerical illustrations matching the predicted convergence rates of Theorems 4.2 and 4.3 are provided in [25].

To compare with Theorem 2.4, we take $r = s$ and assume that Ω is convex, which allows for any α in $(0, 1]$. We choose a number of sinc quadrature points $N_+ \approx N_- \approx \log(1/h)$ so that sinc quadrature and finite element errors are balanced. Therefore, Theorems 4.2 and 4.3 yield for the Dunford-Taylor method

$$\|u - U^k\|_{\mathbb{H}^s(\Omega)} \lesssim (\#\mathcal{T})^{-2\alpha_*/d} \|f\|_{\mathbb{H}^\sigma(\Omega)},$$

where, after the slight modification leading to (4.5), $\sigma := \max(2\alpha_* - s, 0)$ and $(\#\mathcal{T})^{-1/d} \approx h$ for quasi-uniform subdivisions, provided we discard logarithmic terms. Section 2.7 and [18, 88] show that the PDE approach introduced in [95] yields the following error estimate

$$\|u - U\|_{\mathbb{H}^s(\Omega)} \lesssim (\#\mathcal{T})^{-1/d} \|f\|_{\mathbb{H}^{1-s}(\Omega)},$$

when a hp -FEM is used in the extended variable y after discarding logarithmic terms; $\#\mathcal{T}$ corresponds to the number of degrees of freedom. This estimate exhibits quasi-optimal linear order for the regularity $f \in \mathbb{H}^{1-s}(\Omega)$. We also see that the Dunford-Taylor method possesses the optimal rate of convergence $2\alpha_* = 2 - s > 1$ allowed by polynomial interpolation theory for smoother datum $f \in \mathbb{H}^\sigma(\Omega)$, $\sigma = 2(1 - s)$. It is worth noting as well that the same regularity on f is required by the Dunford-Taylor method to deliver the linear order of convergence exhibited by the extension method.

The Dunford-Taylor algorithm seems advantageous in a multi-processor context as it appears to exhibit good strong and weak scaling properties. The former consists of increasing the number of processors for a fixed number of degrees of freedom, while for the latter, the number of degrees of freedom per processor is kept constant when increasing the problem size. We refer to [47] for a comparison of these methods and others not discussed in this overview. However, it is important to mention that this reference does not take into consideration newer developments for the method of Section 2, like the diagonalization procedure discussed in Section 2.7. In fact, the diagonalization approach reduces the full tensor product approach of a hp -FEM on the extended variable y with a P_1 -FEM in Ω of [18] to a sequence of decoupled parallelizable singularly perturbed problems in Ω : $\log(\#\mathcal{T}_\Omega)$ problems of size $\#\mathcal{T}_\Omega$. In this case, the extended method can also be efficiently implemented in a multi-processor context and should be subject to further studies.

Finally, we mention that in [18], several additional improvements of the extended method are provided. Whether these improvements can be transferred to the Dunford-Taylor method remains open.

4.2.1. Numerical Experiment. We consider the unit square $\Omega = (0, 1)^2 \subset \mathbb{R}^2$ and study numerically the efficiency in the approximation of u the solution of

$$(-\Delta)^s u = f, \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega$$

where $f : \Omega \rightarrow \mathbb{R}$ is given for all $(x_1, x_2) \in \Omega$ by

$$(4.6) \quad f(x_1, x_2) = \begin{cases} 1 & \text{if } (x_1 - 0.5)(x_2 - 0.5) > 0 \\ 0 & \text{otherwise.} \end{cases}$$

The convergence rate predicted by Theorem 4.2 when measuring the errors $L^2(\Omega)$ ($r = 0$) depends on the regularity of f and the pick-up regularity index α . Since Ω is convex, the pick-up regularity index α can be chosen to be any number in $(0, 1]$ but the error decay rate is restricted by the regularity of f . In this context $f \in \mathbb{H}^\delta(\Omega)$ with $\delta = \frac{1}{2} - \epsilon$ for all $\epsilon > 0$ and condition $\delta \geq \gamma = \max(2(\alpha_* - s), 0)$




FIGURE 4. Approximations of $(-\Delta)^{-s}f$ for the checkerboard righthand side (4.6) and for $s = 0.1$ (left). The regularization effect of s on the solution is highlighted in the right plot depicting the approximate solutions for different values of s along the line $\{y = 0.25\}$ ($s = 0.1$ dashed line, $s = 0.5$ doted line, $s = 0.8$ solid line).

holds provided $\alpha_* = \alpha < s + \frac{1}{4}$. Hence, discarding the sinc quadrature error, the predicted error decay is proportional to h^β (up to a log factor), with

$$\beta = \begin{cases} 2, & \text{if } s > \frac{3}{4}, \\ 2(s + \frac{1}{4}), & \text{otherwise.} \end{cases}$$

The errors $\|u - u_h^N\|_{L_2(\Omega)}$ are computed using the first 300 modes in x and y (90000 modes) of the Fourier representation of the exact solution and the number of quadrature points is taken large enough not to influence the space discretization error. The average rate of convergence $AROC$ is given by the average of observed rates of convergence

$$\ln(e_i/e_{i+1}) / \ln(h_i/h_{i+1}), \quad i = 1, 2, 3,$$

computed using four meshes and where e_i are the L_2 errors and h_i the diameter of the quasi-uniform subdivision i . We report the $AROC$ in Table 3, which are comparable with the rate $h^{2\beta}$ predicted by Theorem 4.2.

s	$s < \frac{3}{4}$							$s > \frac{3}{4}$	
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
AROC	0.92	1.06	1.22	1.4	1.52	1.72	1.86	1.94	1.96
THM	0.7	0.9	1.1	1.3	1.5	1.7	1.9	2.0	2.0

TABLE 3. Average observed rate of convergence (AROC) for different values of s compared with the rates predicted by Theorem 4.2 (THM). The average observed rates of convergence are consistent with the predicted rates in all cases.

The approximate solution for $s = 0.1$ is provided in Figure 4 together with a cut over the horizontal line $\{y = 0.25\}$ of the approximate solutions corresponding to $s = 0.1, 0.5, 0.8$, thereby illustrating the regularizing effect of s .

Additional numerical observations can be found in [25, 26].

4.2.2. *Extensions.* We now discuss several extensions.

- *Symmetric operators and other boundary conditions.* The operator $L = -\Delta$ can be replaced by any symmetric elliptic operators as long as the associated bilinear form $(u, w) \mapsto \int_{\Omega} Lu w$ remains coercive and bounded in $H_0^1(\Omega)$. Different boundary conditions can be considered similarly as well. However, it is worth pointing out that the characterization of $\mathcal{D}(L^r)$ depends on the boundary condition and must be established.

As an illustration, Figure 5 depicts the approximations using parametric surface finite element [58] of the solution to

$$(4.7) \quad (-\Delta_{\Gamma})^s u = 1 \quad \text{on } \Gamma, \quad u = 0 \quad \text{on } \partial\Gamma,$$

where Δ_{Γ} is the surface Laplacian on $\Gamma \subset \mathbb{R}^3$, either the side boundary of a cylinder or given by

$$(4.8) \quad \Gamma := \{(x_1 + 2 \sin(x_3), x_2 + 2 \cos(x_3), 10x_3) \in \mathbb{R}^3 : (x_1, x_2, x_3) \in \mathbb{S}_2, x_3 \geq 0\}.$$




FIGURE 5. Dunford-Taylor Method: Numerical approximation of the solution to the spectral Laplacian problem (4.7) on an hyper-surface (darker = smaller values; lighter = larger values). (Left) $s = 0.8$ and Γ is the side boundary of a cylinder of radius 1 and height 2; (Right) $s = 0.5$ and Γ is given by (4.8).

- *Regularly accretive operators.* The class of operators L can be extended further to a subclass of non-symmetric operators. They are the unbounded operators associated with coercive and bounded sesquilinear forms in $H_0^1(\Omega)$ (regularly accretive operators [73]). In this case, fractional powers cannot be defined using a spectral decomposition as in (1.6) but rather directly using the Dunford-Taylor representation (1.10) and the Balakrishnan formula (1.11), which remain valid. The bottleneck is the characterization of the functional spaces $\mathcal{D}(L^{r/2})$ in terms of Sobolev regularity. It turns out that for $-1 < r < 1$, we have that $\mathcal{D}(L^{r/2})$ is the same as for the symmetric operator [72]

$$\mathcal{D}(L^{r/2}) = \mathcal{D}((L + L^*)^{r/2}) = \mathbb{H}^r(\Omega),$$

where L^* denotes the adjoint of L . This characterization does not generally hold for $r = 1$ (Kato square root problem [73]). McKintosh [86] proved that $\mathcal{D}(L^{1/2}) = \mathbb{H}^1(\Omega) = H_0^1(\Omega)$ for sesquilinear forms of the type

$$(\psi, \varphi) \mapsto \int_{\Omega} (B \nabla \psi \cdot \nabla \bar{\varphi} + \beta_1 \cdot \nabla \psi \bar{\varphi} + \psi \beta_2 \cdot \nabla \bar{\varphi} + c \psi \bar{\varphi}) dx,$$

where $B \in L^\infty(\Omega, \text{GL}(\mathbb{R}^d))$, $\beta_1, \beta_2 \in L^\infty(\Omega, \mathbb{R}^d)$ and $c \in L^\infty(\Omega)$ are such that the form is coercive and bounded. This characterization is extended in [26,

Theorem 6.4] up to $r = 1 + \alpha$ so that similar convergence estimates to those in Theorems 4.2 and 4.3 are established. To illustrate the method for non-symmetric operators, we consider the following example

$$\left(-\Delta + \begin{pmatrix} 1 \\ 1 \end{pmatrix} \cdot \nabla\right)^s u = 1, \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

where $\Omega = (-1, 1)^2 \setminus [0, 1]^2$ is a L-shaped domain. Figure 6 reports the fully discrete approximations given by (4.4) for $s = 0.2, 0.5, 0.8$.




FIGURE 6. Dunford-Taylor Method: Approximation of the solution to fractional advection - diffusion problem on a L-shaped domain. (Left) Solution with isolines for $s = 0.5$. (Right) Plots of u for $s = 0.2, 0.5, 0.8$ on the segment from the corner opposite to the re-entrant corner (of coordinate $(-1, 1)$) to the re-entrant corner (of coordinate $(0, 0)$). It appears that the boundary layer intensity (but not its width) at the re-entrant corner depends on the power fraction s .

- *Space and time fractional diffusion.* In [23, 22] the space-time fractional problem

$$\partial_t^\gamma u + (-\Delta)^s u = f, \quad u(0) = u_0,$$

is studied, where ∂_t^γ denotes the so-called Caputo derivative of order $\gamma \in (0, 1]$; see (2.18). The solution of the space-time fractional problem is given by [111]

$$u(t) = e_{\gamma,1}(-t^\gamma L^s)u_0 + \int_0^t \xi^{\gamma-1} e_{\gamma,\gamma}(-\xi^\gamma L^s) f(t-\xi) d\xi,$$

where, again in this case, a Dunford-Taylor representation can be used to write

$$e_{\gamma,\mu}(-t^\gamma L^s) = \frac{1}{2\pi i} \int_{\mathcal{C}} e_{\gamma,\mu}(-t^\gamma z^s) (z - L)^{-1} dz$$

and $e_{\gamma,\mu}$ defined on \mathbb{C} is the Mittag-Leffler function. Because of the presence of $e_{\gamma,\mu}$, the contour \mathcal{C} cannot be deformed onto the negative real axis anymore, which prevents a representation like in (1.11). Instead, the sinc quadrature is performed directly on a hyperbolic parametrization of \mathcal{C} in the complex plane. Nevertheless, we obtain the error estimates

$$\|u(t) - U^k(t)\|_{L^2(\Omega)} \lesssim \left(t^{-\gamma\alpha_*/s} h^{2\alpha_*} + t^{-\gamma} e^{-c/k} \right) \|u_0\|_{L^2(\Omega)},$$

when $f = 0$ and where $\alpha_* := \min(\alpha, s^-)$ with s^- denoting any number strictly smaller than s . Here c is a constant independent of h and k . We refer the reader to [23, 22] for estimates measuring the error in higher norms or for improved results (in the singularity when $t \rightarrow 0$) when u_0 is smoother. Note that the representation used does not need a time-stepping method for the initial value problem.

When $u_0 = 0$ instead, a graded (a-priori known and depending on γ) mesh in time towards $t = 0$ is put forward. A midpoint quadrature scheme (second order) in time for a total of $\mathcal{N} \log(\mathcal{N})$ time steps yields the error estimates

$$\begin{aligned} \|u(t) - U^{k,\mathcal{N}}(t)\|_{L^2(\Omega)} &\lesssim t^{(1-\alpha_*/s)\gamma} h^{2\alpha_*} \|f\|_{L^2((0,t)\times\Omega)} \\ &\quad + \max\{t^\gamma, t^{3/2+\gamma}\} \mathcal{N}^{-2} \|f\|_{H^2(0,t;L^2(\Omega))} \\ &\quad + \log(\mathcal{N}) e^{-c/k} \|f\|_{L^\infty(0,t;L^2(\Omega))}. \end{aligned}$$

Notice that the method exhibits *second order* convergence rate (up to a logarithmic term) with respect to the number of time intervals. Again, we refer to [22, 23] for more details as well as additional estimates when measuring the error in higher order norms.

4.3. Integral Laplacian. The strategy used for the spectral Laplacian in the previous section cannot be used for the integral Laplacian. In fact, formulas like (1.10) are not well defined.

Instead, we rely on the following equivalent representation of the bilinear form (4.1) in the weak formulation (1.12). Recall that fractional order Sobolev spaces in \mathbb{R}^d are defined and normed by

$$H^r(\mathbb{R}^d) = \left\{ w \in L^2(\mathbb{R}^d) : \|w\|_{H^r(\mathbb{R}^d)} = \left(\int_{\mathbb{R}^d} (1 + |\xi|^2)^{r/2} |\mathcal{F}(w)(\xi)|^2 d\xi \right)^{1/2} < \infty \right\}$$

for $r > 0$, and that the notation \tilde{w} stands for the zero extension of w outside Ω so that $w \in \mathbb{H}^r(\Omega)$ if and only if $\tilde{w} \in H^r(\mathbb{R}^d)$ for $r \in [0, 3/2)$; see definition (1.5).

Theorem 4.5 (equivalent representation). *Let $s \in (0, 1)$ and $0 \leq r \leq s$. For $\eta \in H^{r+s}(\mathbb{R}^d)$ and $\theta \in H^{s-r}(\mathbb{R}^d)$,*

$$\begin{aligned} &\int_{\mathbb{R}^d} |\xi|^{r+s} \mathcal{F}(\eta)(\xi) |\xi|^{s-r} \overline{\mathcal{F}(\theta)(\xi)} d\xi \\ &= \frac{2 \sin(s\pi)}{\pi} \int_0^\infty \mu^{1-2s} \int_{\mathbb{R}^d} ((-\Delta)(I - \mu^2 \Delta)^{-1} \eta) \theta dx d\mu. \end{aligned}$$

To prove the above theorem, it suffices to note that using Parseval's theorem

$$\int_{\mathbb{R}^d} ((-\Delta)(I - \mu^2 \Delta)^{-1} \eta) \theta dx = \int_{\mathbb{R}^d} \frac{|\xi|^2}{1 + \mu^2 |\xi|^2} \mathcal{F}(\eta)(\xi) \overline{\mathcal{F}(\theta)(\xi)} d\xi.$$

and use the change of variable $t = \mu|\xi|$ together with the relation

$$\int_0^\infty \frac{t^{1-2s}}{1+t^2} dt = \frac{\pi}{2 \sin(\pi s)}.$$

For more details, we refer to [24, Theorem 4.1].

In order to make the above representation more amenable to numerical methods, for $\psi \in L^2(\mathbb{R}^d)$, we define $v(\psi, \mu) := v(\mu) \in H^1(\mathbb{R}^d)$ to be the solution to

$$(4.9) \quad \int_{\mathbb{R}^d} v(\mu) \phi dx + \mu^2 \int_{\mathbb{R}^d} \nabla v(\mu) \cdot \nabla \phi dx = - \int_{\mathbb{R}^d} \psi \phi dx, \quad \forall \phi \in H^1(\mathbb{R}^d).$$

Using this notation along with definition (1.5) of $\mathbb{H}^s(\Omega)$, we realize that for $\eta, \theta \in \mathbb{H}^s(\Omega)$ with $s \in (0, 1)$, we have

$$\int_{\mathbb{R}^d} |\xi|^s \mathcal{F}(\tilde{\eta})(\xi) |\xi|^s \overline{\mathcal{F}(\tilde{\theta})(\xi)} d\xi = \frac{2 \sin(s\pi)}{\pi} \int_0^\infty \mu^{-1-2s} \left(\int_{\Omega} (\eta + v(\tilde{\eta}, \mu)) \theta dx \right) d\mu;$$

note that $v(\tilde{\eta}, \mu)$ does not vanish outside Ω . This prompts the definition

$$(4.10) \quad a(\eta, \theta) := \frac{2 \sin(s\pi)}{\pi} \int_0^\infty \mu^{-1-2s} \left(\int_{\Omega} (\eta + v(\tilde{\eta}, \mu)) \theta dx \right) d\mu,$$

for $\eta, \theta \in \mathbb{H}^s(\Omega)$. The above representation is the starting point of the proposed numerical method. The solution $u \in \mathbb{H}^s(\Omega)$ of the fractional Laplacian satisfies

$$(4.11) \quad a(u, w) = \langle f, w \rangle \quad \forall w \in \mathbb{H}^s(\Omega).$$

We discuss in Section 4.3.4 a Strang's type argument to assess the discretization error from the consistency errors generated by the approximation of $a(\cdot, \cdot)$ using sinc quadratures, domain truncations, and finite element discretizations.

4.3.1. *Sinc Quadrature.* We proceed as in Section 4.2 for the spectral Laplacian and use the change of variable $\mu = e^{-\frac{1}{2}y}$ to arrive at

$$a(\eta, \theta) = \frac{\sin(s\pi)}{\pi} \int_{-\infty}^{\infty} e^{sy} \left(\int_{\Omega} (\eta + v(\tilde{\eta}, \mu(y))) \theta \, dx \right) dy.$$

Given a quadrature spacing $k > 0$, N_+ and N_- two positive integers, the sinc quadrature approximation of $a(\cdot, \cdot)$ is given by

$$(4.12) \quad a^k(\eta, \theta) := \frac{\sin(s\pi)}{\pi} k \sum_{\ell=-N_-}^{N_+} e^{sy_\ell} \int_{\Omega} (\eta + v(\tilde{\eta}, \mu(y_\ell))) \theta \, dx.$$

Notice that we only emphasize the dependency in k in the approximation of $a(\cdot, \cdot)$ as we will select N_+ and N_- as a function of k .

The consistency error between $a^k(\cdot, \cdot)$ and $a(\cdot, \cdot)$ is described in the following result. We simply note that, as for the spectral Laplacian discussed in Section 4.1, the proof of Theorem 4.6 is given in [24, Theorem 5.1 and Remark 5.1] and relies on the holomorphic property and decay as $\mu \rightarrow \infty$ of the integrand in (4.10).

Theorem 4.6 (quadrature consistency). *Given $\theta \in \mathbb{H}^s(\Omega)$ and $\eta \in \mathbb{H}^\delta(\Omega)$ with $s < \delta \leq \min(2-s, \sigma)$, where σ stands for any number strictly smaller than $3/2$. Set $N_+ := \lceil \frac{\pi^2}{2k^2(\delta-s)} \rceil$ and $N_- := \lceil \frac{\pi^2}{4sk^2} \rceil$. Then, we have*

$$|a(\eta, \theta) - a^k(\eta, \theta)| \lesssim \max\left(\frac{1}{\delta-s}, \frac{1}{s}\right) e^{-\pi^2/(4k)} \|\eta\|_{\mathbb{H}^\delta(\Omega)} \|\theta\|_{\mathbb{H}^s(\Omega)}.$$

4.3.2. *Truncated Problems.* The sinc approximation of $a(\cdot, \cdot)$ defined by (4.12) requires the computation of $v(\tilde{\eta}, \mu(y_\ell))$ for each quadrature point y_ℓ (here $\eta \in \mathbb{H}^\beta(\Omega)$ for some $s < \beta < 3/2$ is fixed). This necessitates, according to (4.9), the approximations of $(I - \mu(y_\ell)^2 \Delta)^{-1}$ on \mathbb{R}^d . The proposed method relies on truncations of this infinite domain problem and uses standard finite elements on the resulting bounded domains. As we shall see, the truncated domain diameter must depend on the quadrature point y_ℓ .

We let B be a convex bounded domain containing Ω and the origin of \mathbb{R}^d . Without loss of generality, we assume that the diameter of B is 1. For a truncation parameter M , we define the dilated domains

$$(4.13) \quad B^M(\mu) := \begin{cases} \{y = (1 + \mu(1 + M))x : x \in B\}, & \mu \geq 1, \\ \{y = (2 + M)x : x \in B\}, & \mu < 1, \end{cases}$$

and for $\psi \in L^2(\mathbb{R}^d)$, the associated functions $v^M(\mu) := v^M(\psi, \mu) \in H_0^1(B^M(\mu))$ satisfying

$$(4.14) \quad \begin{aligned} \int_{B^M(\mu)} v^M(\mu) w \, dx + \mu^2 \int_{B^M(\mu)} \nabla v^M(\mu) \cdot \nabla w \, dx \\ = - \int_{B^M(\mu)} \psi w \, dx \quad \forall w \in H_0^1(B^M(\mu)); \end{aligned}$$

compare with (4.9). The exponential decay of the function $v(\tilde{\eta}, \mu)$ yields

$$\|v(\tilde{\eta}, \mu) - v^M(\tilde{\eta}, \mu)\|_{L^2(B^M(\mu))} \lesssim e^{-\max(1, \mu)cM/\mu} \|\eta\|_{L^2(\Omega)},$$

where c is a constant independent of M and μ (see [24, Lemma 6.1]). As a consequence, the truncation consistency in using

$$a^{k,M}(\eta, \theta) := \frac{\sin(s\pi)}{\pi} k \sum_{\ell=-N_-}^{N_+} e^{sy_\ell} \int_{\Omega} (\eta + v^M(\tilde{\eta}, \mu(y_\ell))) \theta \, dx$$

instead of $a^k(\eta, \theta)$ decays exponentially fast as a function of M [24, Theorem 6.2].

Theorem 4.7 (truncation consistency). *For M sufficiently large, there is positive constant c independent of M and k such that for all $\eta, \theta \in L^2(\Omega)$*

$$|a^k(\eta, \theta) - a^{k,M}(\eta, \theta)| \lesssim e^{-cM} \|\eta\|_{L^2(\Omega)} \|\theta\|_{L^2(\Omega)}.$$

4.3.3. Finite Element Discretization. We now turn our attention to the finite element approximation of $v^M(\tilde{\eta}, \mu)$ defined by (4.14). For simplicity, we assume that the domain Ω is polytopal so that it can be partitioned into a conforming subdivision \mathcal{T} with elements of maximum diameter h as in Section 4.1.1. Generic constants below may depend on the shape regularity and quasi-uniformity constants of \mathcal{T} without mention of it.

We need two subspaces of globally continuous piecewise linear polynomials. The first one, $\mathbb{U}(\mathcal{T}) \subset H_0^1(\Omega)$, is defined in (2.10) relative to the partition \mathcal{T} . The second subspace, denoted $\mathbb{U}(\mathcal{T}^M(\mu))$, has a similar definition but relative to the subdivision $\mathcal{T}^M(\mu)$ of $B^M(\mu(y_\ell))$. We impose that the partitions $\mathcal{T}^M(\mu)$ match in Ω , which implies that restrictions of functions in $\mathbb{U}(\mathcal{T}^M(\mu))$ are continuous piecewise linears over \mathcal{T} . We refer to Section 4.3.5 and [24] for details on the constructions of such partitions, which is the bottleneck of the proposed method.

We now define for any $\psi \in L^2(\mathbb{R}^d)$ the finite element approximation $V^M(\mu) = V^M(\psi, \mu) \in \mathbb{U}(\mathcal{T}^M(\mu))$ of the function $v^M(\mu)$ to be

$$\begin{aligned} \int_{B^M(\mu)} V^M(\mu) W \, dx + \mu^2 \int_{B^M(\mu)} \nabla V^M(\mu) \cdot \nabla W \, dx \\ = - \int_{B^M(\mu)} \psi W \, dx \quad \forall W \in \mathbb{U}(\mathcal{T}^M(\mu)). \end{aligned}$$

The fully discrete approximation of the bilinear form $a(\cdot, \cdot)$ then reads

$$(4.15) \quad a_{\mathcal{T}}^{k,M}(\Xi, \Theta) := \frac{\sin(s\pi)}{\pi} k \sum_{\ell=-N_-}^{N_+} e^{sy_\ell} \int_{\Omega} (\Xi + V^M(\tilde{\Xi}, \mu(y_\ell))) \Theta \, dx \quad \forall \Xi, \Theta \in \mathbb{U}(\mathcal{T}).$$

Note that $V^M(\tilde{\Xi}, \mu)$ is piecewise linear over \mathcal{T} and the sum $\Xi + V^M(\tilde{\Xi}, \mu)$ is easy to perform. The consistency error between $a^{k,M}(\cdot, \cdot)$ and $a_{\mathcal{T}}^{k,M}(\cdot, \cdot)$ is given next; see [24, Theorem 7.5] for a proof.

Theorem 4.8 (finite element consistency). *For $\beta \in (s, 3/2)$ the following estimate is valid*

$$|a^{k,M}(\Xi, \Theta) - a_{\mathcal{T}}^{k,M}(\Xi, \Theta)| \lesssim (1 + |\log h|) h^{\beta-s} \|\Xi\|_{\mathbb{H}^\beta(\Omega)} \|\Theta\|_{\mathbb{H}^s(\Omega)},$$

for all $\Xi, \Theta \in \mathbb{U}(\mathcal{T})$.

4.3.4. Strang's Lemma. In addition to the three consistency estimates described above, Strang's Lemma requires the $\mathbb{U}(\mathcal{T})$ -ellipticity of the fully discrete form $a_{\mathcal{T}}^{k,M}(\cdot, \cdot)$. To show this, [24] invokes Theorem 4.6 (quadrature consistency) with $\delta := \min\{2 - s, \beta\}$ and an inverse estimate to write for $\Xi, \Theta \in \mathbb{U}(\mathcal{T})$

$$|a^k(\Xi, \Theta) - a(\Xi, \Theta)| \lesssim e^{-\pi^2/(4k)} h^{s-\delta} \|\Xi\|_{\mathbb{H}^s(\Omega)} \|\Theta\|_{\mathbb{H}^s(\Omega)}.$$

This, together with the monotonicity property

$$a_{\mathcal{T}}^{k,M}(\Xi, \Theta) \geq a^k(\Xi, \Theta),$$

and the coercivity of the exact bilinear form $a(\cdot, \cdot)$ in $\mathbb{H}^s(\Omega)$, yields the $\mathbb{U}(\mathcal{T})$ -ellipticity of $a_{\mathcal{T}}^{k,M}(\cdot, \cdot)$ provided

$$(4.16) \quad e^{-\pi^2/(4k)} h^{s-\delta} \leq c$$

for an explicit constant c .

The fully discrete approximation $U^{k,M} \in \mathbb{U}(\mathcal{T})$ of u satisfying (4.11) is given by

$$a_{\mathcal{T}}^{k,M}(U^{k,M}, W) = \int_{\Omega} fW \, dx \quad \forall W \in \mathbb{U}(\mathcal{T}).$$

To measure the discrepancy between u and $U^{k,M}$ in $\mathbb{H}^s(\Omega)$, we assume the additional regularity $u \in \mathbb{H}^{\beta}(\Omega)$ for some $\beta \in (s, 3/2)$. The expected regularity of u , solution to the integral fractional Laplacian, is discussed in Theorems 3.1 and 3.3. The theorem below, proved in [24, Theorem 7.7]), guarantees that the proposed method delivers an *optimal* rate of convergence (up to a logarithmic factor).

Theorem 4.9 (error estimate). *Assume that (4.16) holds and that $u \in \mathbb{H}^{\beta}(\Omega)$ for some $\beta \in (s, 3/2)$. Then there is a constant c independent of h , M , and k such that*

$$\|u - U^{k,M}\|_{\mathbb{H}^s(\Omega)} \lesssim \left(e^{-\pi^2/(4k)} + e^{-cM} + |\log h| h^{\beta-s} \right) \|u\|_{\mathbb{H}^{\beta}(\Omega)}.$$

Remark 4.10 (implementation). Let us denote by \mathbf{U} the coefficient vector of the finite element approximation $U^{k,M}$ and by \mathbf{F} to be the coefficient vector of the L^2 projection of f onto $\mathbb{U}(\mathcal{T})$. The mass and stiffness matrix in $\mathbb{U}(\mathcal{T})$ are denoted by \mathcal{M} and \mathcal{A} while those in $\mathbb{U}(\mathcal{T}^M(\mu))$ by $\mathcal{M}(\mu)$ and $\mathcal{A}(\mu)$. We also denote by $\mathcal{E}(\mu) : \mathbb{R}^{\dim(\mathbb{U}(\mathcal{T}))} \rightarrow \mathbb{R}^{\dim(\mathbb{U}(\mathcal{T}^M(\mu)))}$ the extension by zero operator and by $\mathcal{R}(\mu)$ its dual (restriction) operation. With these notations and the representation (4.15), we realize that \mathbf{U} satisfies

$$(4.17) \quad \frac{\sin(\pi s)k}{\pi} \sum_{\ell=-N_-}^{N_+} e^{sy_{\ell}} \mathcal{M}\mathcal{R}(\mu_{\ell})(e^{y_{\ell}} \mathcal{M}(\mu_{\ell}) + \mathcal{A}(\mu_{\ell}))^{-1} \mathcal{E}(\mu_{\ell}) \mathcal{A} \mathbf{U} = \mathbf{F},$$

where $y_{\ell} = \ell k$ and $\mu_{\ell} = e^{-\frac{1}{2}y_{\ell}}$. A preconditioned conjugate gradient is used in practice to find \mathbf{U} , i.e.,

$$\frac{\sin(\pi s)k}{\pi} \sum_{\ell=-N_-}^{N_+} e^{sy_{\ell}} \mathcal{M}\mathcal{R}(\mu_{\ell})(e^{y_{\ell}} \mathcal{M}(\mu_{\ell}) + \mathcal{A}(\mu_{\ell}))^{-1} \mathcal{E}(\mu_{\ell}) \mathcal{A} \mathbf{V}$$

need to be computed for some given $\mathbf{V} \in \mathbb{R}^{\dim(\mathbb{U}(\mathcal{T}))}$.

Notice that the error estimate stated in Theorem 4.9 for quasi-uniform meshes is of order about $h^{1/2}$ and is similar to the one derived for the integral method in Theorem 3.7. To see this, we choose $\beta = s + \frac{1}{2} - \varepsilon$ with $\varepsilon > 0$ arbitrary, which is consistent with the regularity of u guaranteed by Theorem 3.3, along with $M \approx \log(1/h)$ and $N_+ \approx N_- \approx |\log h|^2$ to balance the three sources of errors. It is worth mentioning that using graded meshes for $d = 2$, Theorem 3.10 states an optimal linear rate of convergence (up to a logarithmic factor) provided $s \in (1/2, 1)$. Whether such a strategy applies to the Dunford-Taylor method remains open.

4.3.5. *Numerical Experiment.* To illustrate the method, we depict in Figure 7 the approximation $U^{k,M}$ for $s = 0.3$, $f = 1$, and $\Omega = B(0, 1)$, the unit ball in \mathbb{R}^3 . In this case, we let B in (4.13) to be Ω . The subdivisions of the computational domains $B^M(\mu(y_\ell))$ at each quadrature points y_ℓ are then generated as follow. We start from a subdivision \mathcal{T} of B and denote by \mathcal{S} the corresponding subdivision of ∂B . We then expand radially \mathcal{S} to obtain several slabs in between B and $B^M(\mu(y_\ell))$. The slabs are connected using radial lines emanating from 0 and passing through all the vertices of \mathcal{S} . We refer to [24, Section 8] for further discussions on mesh generation, matrix representation of the fully discrete scheme, and a preconditioned iterative method.




FIGURE 7. Dunford-Integral Method: Numerical approximation of the solution to the fractional integral Laplacian for $s = 0.3$ and $f = 1$ in the unit ball of \mathbb{R}^3 (darker = 0.0, whiter = 0.7). The lines represent the isosurfaces $\{u(x) = k/10\}$ for $k = 0, \dots, 7$ and reflect the singular behavior of the solution towards the domain boundary, see (1.7).

REFERENCES

- [1] S. Abe and S. Thurner. Anomalous diffusion in view of Einstein’s 1905 theory of Brownian motion. *Physica A: Statistical Mechanics and its Applications*, 356(2–4):403 – 407, 2005.
- [2] M. Abramowitz and I.A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [3] G. Acosta, F. Bersetche, and J.P. Borthagaray. Finite element approximations for fractional evolution problems. *arXiv:1705.09815v1*, 2017.
- [4] G. Acosta, F. M. Bersetche, and J.P. Borthagaray. A short FE implementation for a 2d homogeneous Dirichlet problem of a fractional Laplacian. *Comput. Math. Appl.*, 74(4):784–816, 2017.
- [5] G. Acosta and J.P. Borthagaray. A fractional Laplace equation: regularity of solutions and finite element approximations. *SIAM J. Numer. Anal.*, 55(2):472–495, 2017.
- [6] G. Acosta, J.P. Borthagaray, O. Bruno, and M. Maas. Regularity theory and high order numerical methods for one-dimensional fractional-Laplacian equations. *Math. Comp.*, 2017. Accepted for publication.
- [7] G. Acosta, J.P. Borthagaray, and N. Heuer. Finite element approximations for the nonhomogeneous fractional Dirichlet problem. In preparation.

- [8] M. Ainsworth and C. Glusa. Towards an efficient finite element method for the integral fractional Laplacian on polygonal domains. *arXiv:1708.01923v1*, 2017.
- [9] H. Antil and E. Otárola. A FEM for an optimal control problem of fractional powers of elliptic operators. *SIAM J. Control Optim.*, 53(6):3432–3456, 2015.
- [10] H. Antil and E. Otárola. An a posteriori error analysis for an optimal control problem involving the fractional Laplacian. *IMA J. Numer. Anal.*, 2017. (to appear).
- [11] H. Antil, E. Otárola, and A.J. Salgado. Optimization with respect to order in a fractional diffusion model: analysis, approximation and algorithm aspects. *arXiv:1612.08982v1*, 2016.
- [12] H. Antil, E. Otárola, and A.J. Salgado. A space-time fractional optimal control problem: analysis and discretization. *SIAM J. Control Optim.*, 54(3):1295–1328, 2016.
- [13] I. Babuška and A. Miller. A feedback finite element method with a posteriori error estimation. I. The finite element method and some basic properties of the a posteriori error estimator. *Comput. Methods Appl. Mech. Engrg.*, 61(1):1–40, 1987.
- [14] I. Babuška and J. Osborn. Eigenvalue problems. In *Handbook of numerical analysis, Vol. II*, Handb. Numer. Anal., II, pages 641–787. North-Holland, Amsterdam, 1991.
- [15] C. Bacuta, J.H. Bramble, and J.E. Pasciak. New interpolation results and applications to finite element methods for elliptic boundary value problems. *East-West J. Numer. Math.*, 3:179–198, 2001.
- [16] W. Bangerth, R. Hartmann, and G. Kanschat. deal.II—differential equations analysis library. *Technical Reference: <http://dealii.org>*.
- [17] W. Bangerth, R. Hartmann, and G. Kanschat. deal.II—a general-purpose object-oriented finite element library. *ACM Trans. Math. Software*, 33(4):Art. 24, 27, 2007.
- [18] L. Banjai, J.M. Melenk, R.H. Nochetto, E. Otárola, A.J. Salgado, and Ch. Schwab. Tensor FEM for spectral fractional diffusion. *arXiv:1707.07367v1*, 2017.
- [19] J. Bertoin. *Lévy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- [20] M.Š. Birman and M.Z. Solomjak. *Spektralnaya teoriya samosopryazhennykh operatorov v gilbertovom prostranstve*. Leningrad. Univ., Leningrad, 1980.
- [21] A. Bonito, J.-L. Guermond, and F. Luddens. Regularity of the Maxwell equations in heterogeneous media and Lipschitz domains. *J. Math. Anal. Appl.*, 408(2):498–512, 2013.
- [22] A. Bonito, W. Lei, and J.E. Pasciak. The approximation of parabolic equations involving fractional powers of elliptic operators. *Journal of Computational and Applied Mathematics*, 315:32–48, 2017.
- [23] A. Bonito, W. Lei, and J.E. Pasciak. Numerical approximation of space-time fractional parabolic equations. *Accepted in Comput. Methods Appl. Math.*, 2017.
- [24] A. Bonito, W. Lei, and J.E. Pasciak. Numerical approximation of the integral fractional Laplacian. *arXiv:1707.04290v1*, 2017.
- [25] A. Bonito and J. Pasciak. Numerical approximation of fractional powers of elliptic operators. *Mathematics of Computation*, 84(295):2083–2110, 2015.
- [26] A. Bonito and J.E. Pasciak. Numerical approximation of fractional powers of regularly accretive operators. *IMA Journal of Numerical Analysis*, 2016.
- [27] J.P. Borthagaray and P. Ciarlet, Jr. Nonlocal models for interface problems between dielectrics and metamaterials. In *11th International Congress on Engineered Material Platforms for Novel Wave Phenomena*, 2017.
- [28] J.P. Borthagaray, L.M. Del Pezzo, and S. Martínez. Finite element approximation for the fractional eigenvalue problem. *arXiv:1603.00317v2*, 2017.
- [29] J. Bourgain, H. Brezis, and P. Mironescu. Another look at Sobolev spaces. In *Optimal Control and Partial Differential Equations*, pages 439–455, 2001.
- [30] C. Brändle, E. Colorado, A. de Pablo, and U. Sánchez. A concave-convex elliptic problem involving the fractional Laplacian. *Proc. Roy. Soc. Edinburgh Sect. A*, 143(1):39–71, 2013.
- [31] S.C. Brenner and L.R. Scott. *The mathematical theory of finite element methods*, volume 15 of *Texts in Applied Mathematics*. Springer, New York, third edition, 2008.
- [32] D. Brockmann, L. Hufnagel, and T. Geisel. The scaling laws of human travel. *Nature*, 439(7075):462–465, 2006.
- [33] C. Bucur and E. Valdinoci. *Nonlocal diffusion and applications*, volume 20 of *Lecture Notes of the Unione Matematica Italiana*. Springer; Unione Matematica Italiana, Bologna, 2016.
- [34] X. Cabré and J. Tan. Positive solutions of nonlinear problems involving the square root of the Laplacian. *Adv. Math.*, 224(5):2052–2093, 2010.
- [35] L. Caffarelli and A. Figalli. Regularity of solutions to the parabolic fractional obstacle problem. *J. Reine Angew. Math.*, 680:191–233, 2013.
- [36] L. Caffarelli and Stinga P. Fractional elliptic equations, Caccioppoli estimates, and regularity. *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*, 33:767–807, 2016.

- [37] L. Caffarelli, S. Salsa, and L. Silvestre. Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian. *Invent. Math.*, 171(2):425–461, 2008.
- [38] L. Caffarelli and L. Silvestre. An extension problem related to the fractional Laplacian. *Comm. Part. Diff. Eqs.*, 32(7-9):1245–1260, 2007.
- [39] L. Caffarelli and A. Vasseur. Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation. *Ann. of Math. (2)*, 171(3):1903–1930, 2010.
- [40] A. Capella, J. Dávila, L. Dupaigne, and Y. Sire. Regularity of radial extremal solutions for some non-local semilinear equations. *Comm. Partial Differential Equations*, 36(8):1353–1384, 2011.
- [41] B. Carmichael, H. Babahosseini, S.N. Mahmoodi, and M. Agah. The fractional viscoelastic response of human breast tissue cells. *Physical Biology*, 12(4):046001, 2015.
- [42] P. Carr, H. Geman, D.B. Madan, and M. Yor. The fine structure of asset returns: An empirical investigation. *Journal of Business*, 75:305–332, 2002.
- [43] L. Chen, R.H. Nochetto, E. Otárola, and A.J. Salgado. A PDE approach to fractional diffusion: A posteriori error analysis. *J. Comput. Phys.*, 293:339–358, 2015.
- [44] L. Chen, R.H. Nochetto, E. Otárola, and A.J. Salgado. Multilevel methods for nonuniformly elliptic operators and fractional diffusion. *Math. Comp.*, 85(302):2583–2607, 2016.
- [45] Z.Q. Chen and R. Song. Hardy inequality for censored stable processes. *Tohoku Math. J. (2)*, 55(3):439–450, 2003.
- [46] P. Ciarlet, Jr. Analysis of the Scott-Zhang interpolation in the fractional order Sobolev spaces. *J. Numer. Math.*, 21(3):173–180, 2013.
- [47] R. Čiegis, V. Starikovičius, S. Margenov, and R. Kriauzienė. Parallel solvers for fractional power diffusion problems. *Concurrency and Computation: Practice and Experience*, pages e4216–n/a. e4216 cpe.4216.
- [48] M. Costabel and M. Dauge. General edge asymptotics of solutions of second-order elliptic boundary value problems i. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 123(1):109–155, 1993.
- [49] J. Cushman and T. Glinn. Nonlocal dispersion in media with continuously evolving scales of heterogeneity. *Trans. Porous Media*, 13:123–138, 1993.
- [50] M. Dauge. *Elliptic Boundary Value Problems on Corner Domains*. Lecture Notes in Mathematics, 1341, Springer-Verlag, 1988.
- [51] M. D’Elia and M. Gunzburger. The fractional Laplacian operator on bounded domains as a special case of the nonlocal diffusion operator. *Comput. Math. Appl.*, 66(7):1245 – 1260, 2013.
- [52] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.*, 136(5):521–573, 2012.
- [53] S. Dipierro, X. Ros-Oton, and E. Valdinoci. Nonlocal problems with Neumann boundary conditions. *Rev. Mat. Iberoam.*, 33(2):377–416, 2017.
- [54] J. Duoandikoetxea. *Fourier analysis*, volume 29 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Translated and revised from the 1995 Spanish original by David Cruz-Uribe.
- [55] R.G. Durán and A.L. Lombardi. Error estimates on anisotropic Q_1 elements for functions in weighted Sobolev spaces. *Math. Comp.*, 74(252):1679–1706 (electronic), 2005.
- [56] B. Dyda. A fractional order Hardy inequality. *Illinois J. Math.*, 48(2):575–588, 2004.
- [57] B. Dyda, A. Kuznetsov, and M. Kwaśnicki. Eigenvalues of the fractional Laplace operator in the unit ball. *J. Lond. Math. Soc.*, 95(2):500–518, 2017.
- [58] G. Dziuk. Finite elements for the Beltrami operator on arbitrary surfaces. In *Partial differential equations and calculus of variations*, volume 1357 of *Lecture Notes in Math.*, pages 142–155. Springer, Berlin, 1988.
- [59] A. Einstein. *Investigations on the theory of the Brownian movement*. Dover Publications, Inc., New York, 1956. Edited with notes by R. Fürth, Translated by A. D. Cowper.
- [60] E.B. Fabes, C.E. Kenig, and R.P. Serapioni. The local regularity of solutions of degenerate elliptic equations. *Comm. Part. Diff. Eqs.*, 7(1):77–116, 1982.
- [61] B. Faermann. Localization of the Aronszajn-Slobodeckij norm and application to adaptive boundary element methods. II. The three-dimensional case. *Numer. Math.*, 92(3):467–499, 2002.
- [62] R.K. Gettoor. First passage times for symmetric stable processes in space. *Trans. Amer. Math. Soc.*, 101:75–90, 1961.
- [63] V. Gol’dshhtein and A. Ukhlov. Weighted Sobolev spaces and embedding theorems. *Trans. Amer. Math. Soc.*, 361(7):3829–3850, 2009.
- [64] R. Gorenflo, A.A. Kilbas, F. Mainardi, and S.V. Rogosin. *Mittag-Leffler functions, related topics and applications*. Springer Monographs in Mathematics. Springer, Heidelberg, 2014.

- [65] P. Grisvard. *Elliptic problems in nonsmooth domains*, volume 69 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011. Reprint of the 1985 original [MR0775683], With a foreword by Susanne C. Brenner.
- [66] G. Grubb. Fractional Laplacians on domains, a development of Hörmander’s theory of μ -transmission pseudodifferential operators. *Adv. Math.*, 268:478 – 528, 2015.
- [67] G. Grubb. Spectral results for mixed problems and fractional elliptic operators. *J. Math. Anal. Appl.*, 421(2):1616–1634, 2015.
- [68] L. Hörmander. Ch. II, Boundary problems for “classical” pseudo-differential operators. Available at <http://www.math.ku.dk/~grubb/LH65.pdf>, 1965.
- [69] Y. Huang and A.M. Oberman. Numerical methods for the fractional Laplacian: A finite difference-quadrature approach. *SIAM J. Numer. Anal.*, 52(6):3056–3084, 2014.
- [70] B. Jin, R. Lazarov, and Z. Zhou. Two fully discrete schemes for fractional diffusion and diffusion-wave equations with nonsmooth data. *SIAM J. Sci. Comput.*, 38(1):A146–A170, 2016.
- [71] F. Jochmann. An H^s -regularity result for the gradient of solutions to elliptic equations with mixed boundary conditions. *J. Math. Anal. Appl.*, 238:429–450, 1999.
- [72] T. Kato. Note on fractional powers of linear operators. *Proc. Japan Acad.*, 36:94–96, 1960.
- [73] T. Kato. Fractional powers of dissipative operators. *J. Math. Soc. Japan*, 13:246–274, 1961.
- [74] R.B. Kellogg. Interpolation between subspaces of a Hilbert space. Technical report, Univ. of Maryland,, Inst. Fluid Dynamics and App. Math., Tech. Note BN-719, 1971.
- [75] T. Kilpeläinen. Weighted Sobolev spaces and capacity. *Ann. Acad. Sci. Fenn. Ser. AI Math*, 19(1):95–113, 1994.
- [76] M.A. Krasnosel’skiĭ and Ja.B. Rutickiĭ. *Convex functions and Orlicz spaces*. Translated from the first Russian edition by Leo F. Boron. P. Noordhoff Ltd., Groningen, 1961.
- [77] A. Kufner. *Weighted Sobolev spaces*. A Wiley-Interscience Publication. John Wiley & Sons Inc., New York, 1985. Translated from the Czech.
- [78] A. Kufner and B. Opic. How to define reasonably weighted Sobolev spaces. *Comment. Math. Univ. Carolin.*, 25(3):537–554, 1984.
- [79] A. Kyprianou, A. Osojnik, and T. Shardlow. Unbiased walk-on-spheres’ Monte Carlo methods for the fractional Laplacian. *arXiv:1609.03127v3*, 2017.
- [80] N.S. Landkof. *Foundations of modern potential theory*. Springer-Verlag, New York-Heidelberg, 1972. Translated from the Russian by A. P. Doohovskoy, Die Grundlehren der mathematischen Wissenschaften, Band 180.
- [81] N. Laskin. Fractional quantum mechanics and Lévy path integrals. *Physics Letters A*, 268(4):298–305, 2000.
- [82] C. Lubich. Convolution quadrature and discretized operational calculus. I. *Numer. Math.*, 52(2):129–145, 1988.
- [83] A. Lunardi. *Interpolation theory*. Appunti. Scuola Normale Superiore di Pisa (Nuova Serie). [Lecture Notes. Scuola Normale Superiore di Pisa (New Series)]. Edizioni della Normale, Pisa, second edition, 2009.
- [84] J. Lund and K.L. Bowers. *Sinc methods for quadrature and differential equations*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1992.
- [85] B.M. McCay and M.N.L. Narasimhan. Theory of nonlocal electromagnetic fluids. *Arch. Mech. (Arch. Mech. Stos.)*, 33(3):365–384, 1981.
- [86] A. McIntosh. The square root problem for elliptic operators: a survey. In *Functional-analytic methods for partial differential equations (Tokyo, 1989)*, volume 1450 of *Lecture Notes in Math.*, pages 122–140. Springer, Berlin, 1990.
- [87] W. McLean. *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, Cambridge, 2000.
- [88] D. Meidner, J. Pfefferer, K. Schürholz, and B. Vexler. hp -finite elements for fractional diffusion. *arXiv:1706.04066v1*, 2017.
- [89] R. Metzler and J. Klafter. The restaurant at the end of the random walk: recent developments in the description of anomalous transport by fractional dynamics. *J. Phys. A*, 37(31):R161–R208, 2004.
- [90] P. Morin, R.H. Nochetto, and K.G. Siebert. Local problems on stars: a posteriori error estimators, convergence, and performance. *Math. Comp.*, 72(243):1067–1097 (electronic), 2003.
- [91] B. Muckenhoupt. Weighted norm inequalities for the Hardy maximal function. *Trans. Amer. Math. Soc.*, 165:207–226, 1972.
- [92] R. Musina and A.I. Nazarov. On fractional Laplacians. *Comm. Partial Differential Equations*, 39(9):1780–1790, 2014.
- [93] S. Nazarov and B. Plamenevsky. *Elliptic problems in domains with piecewise smooth boundaries*. De Gruyter expositions in mathematics, De Gruyter, 1994.

- [94] R.H. Nochetto, E. Otárola, and A.J. Salgado. Convergence rates for the classical, thin and fractional elliptic obstacle problems. *Philos. Trans. Roy. Soc. A*, 373(2050):20140449, 14, 2015.
- [95] R.H. Nochetto, E. Otárola, and A.J. Salgado. A PDE approach to fractional diffusion in general domains: a priori error analysis. *Found. Comput. Math.*, 15(3):733–791, 2015.
- [96] R.H. Nochetto, E. Otárola, and A.J. Salgado. A PDE approach to numerical fractional diffusion. In *Proceedings of the 8th International Congress on Industrial and Applied Mathematics*, pages 211–236. Higher Ed. Press, Beijing, 2015.
- [97] R.H. Nochetto, E. Otárola, and A.J. Salgado. A PDE approach to space-time fractional parabolic problems. *SIAM J. Numer. Anal.*, 54(2):848–873, 2016.
- [98] R.H. Nochetto, E. Otárola, and A.J. Salgado. Piecewise polynomial interpolation in Muckenhoupt weighted Sobolev spaces and applications. *Numer. Math.*, 132(1):85–130, 2016.
- [99] R.H. Nochetto, K.G. Siebert, and A. Veerer. Theory of adaptive finite element methods: an introduction. In *Multiscale, nonlinear and adaptive approximation*, pages 409–542. Springer, Berlin, 2009.
- [100] R.H. Nochetto and A. Veerer. Primer of adaptive finite element methods. In *Multiscale and Adaptivity: Modeling, Numerics and Applications, CIME Lectures*. Springer, 2011.
- [101] R.H. Nochetto, T. von Petersdorff, and C.-S. Zhang. A posteriori error analysis for a class of integral equations and variational inequalities. *Numer. Math.*, 116(3):519–552, 2010.
- [102] F.W.J. Olver. *Asymptotics and special functions*. AKP Classics. A K Peters, Ltd., Wellesley, MA, 1997. Reprint of the 1974 original [Academic Press, New York; MR0435697 (55 #8655)].
- [103] E. Otárola. *A PDE approach to numerical fractional diffusion*. ProQuest LLC, Ann Arbor, MI, 2014. Thesis (Ph.D.)—University of Maryland, College Park.
- [104] E. Otárola. A piecewise linear FEM for an optimal control problem of fractional operators: error analysis on curved domains. *ESAIM Math. Model. Numer. Anal.*, 51(4):1473–1500, 2017.
- [105] E. Otárola and A.J. Salgado. Finite element approximation of the parabolic fractional obstacle problem. *SIAM J. Numer. Anal.*, 54(4):2619–2639, 2016.
- [106] E. Otárola and A.J. Salgado. Regularity of solutions to space–time fractional wave equations: a PDE approach. 2017. In preparation.
- [107] E. Otárola and A.J. Salgado. Sparse optimal control for fractional diffusion. *Comput. Math. Appl. Math.*, 2017. (to appear).
- [108] X. Ros-Oton. Nonlocal elliptic equations in bounded domains: a survey. *Publ. Mat.*, 60(1):3–26, 2016.
- [109] X. Ros-Oton and J. Serra. The Dirichlet problem for the fractional Laplacian: regularity up to the boundary. *J. Math. Pures Appl.*, 101(3):275 – 302, 2014.
- [110] X. Ros-Oton and J. Serra. Local integration by parts and Pohozaev identities for higher order fractional Laplacians. *Discrete Contin. Dyn. Syst.*, 35(5):2131–2150, 2015.
- [111] K. Sakamoto and M. Yamamoto. Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *Journal of Mathematical Analysis and Applications*, 382(1):426–447, 2011.
- [112] S.G. Samko, A.A. Kilbas, and O.I. Marichev. *Fractional integrals and derivatives*. Gordon and Breach Science Publishers, Yverdon, 1993. Theory and applications, Edited and with a foreword by S. M. Nikol’skiĭ, Translated from the 1987 Russian original, Revised by the authors.
- [113] S.A. Sauter and C. Schwab. *Boundary element methods*, volume 39 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 2011. Translated and expanded from the 2004 German original.
- [114] L.R. Scott and S. Zhang. Finite element interpolation of nonsmooth functions satisfying boundary conditions. *Math. Comp.*, 54(190):483–493, 1990.
- [115] R. Servadei and E. Valdinoci. On the spectrum of two different fractional operators. *Proc. Roy. Soc. Edinburgh Sect. A*, 144(4):831–855, 2014.
- [116] L. Silvestre. Regularity of the obstacle problem for a fractional power of the Laplace operator. *Comm. Pure Appl. Math.*, 60(1):67–112, 2007.
- [117] D. Sims, E. Southall, N. Humphries, G. Hays, C. Bradshaw, J. Pitchford, A. James, M. Ahmed, A. Brierley, M. Hindell, D. Morritt, M. Musyl, D. Righton, E. Shepard, V. Wearmouth, R. Wilson, M. Witt, and J. Metcalfe. Scaling laws of marine predator search behaviour. *Nature*, 451(7182):1098–1102, 2008.
- [118] J. Sprekels and E. Valdinoci. A new type of identification problems: optimizing the fractional order in a nonlocal evolution equation. *SIAM J. Control Optim.*, 55(1):70–93, 2017.
- [119] P.R. Stinga and J.L. Torrea. Extension problem and Harnack’s inequality for some fractional operators. *Comm. Partial Differential Equations*, 35(11):2092–2122, 2010.

- [120] L. Tartar. *An introduction to Sobolev spaces and interpolation spaces*, volume 3 of *Lecture Notes of the Unione Matematica Italiana*. Springer, Berlin, 2007.
- [121] M.E. Taylor. *Pseudodifferential operators*. Princeton Mathematical Series, vol. 34., 1981.
- [122] B.O. Turesson. *Nonlinear potential theory and weighted Sobolev spaces*, volume 1736 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2000.
- [123] M.I. Višik and G.I. Èskin. Elliptic convolution equations in a bounded region and their applications. *Uspehi Mat. Nauk*, 22(1 (133)):15–76, 1967.
- [124] K. Yosida. *Functional analysis*. Second edition. Die Grundlehren der mathematischen Wissenschaften, Band 123. Springer-Verlag New York Inc., New York, 1968.

(A. Bonito) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA

E-mail address: bonito@math.tamu.edu

(J.P. Borthagaray) IMAS - CONICET AND DEPARTAMENTO DE MATEMÁTICA, FCEYN - UNIVERSIDAD DE BUENOS AIRES, BUENOS AIRES, ARGENTINA

E-mail address: jpbortha@dm.uba.ar

(R.H. Nochetto) DEPARTMENT OF MATHEMATICS AND INSTITUTE FOR PHYSICAL SCIENCE AND TECHNOLOGY, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742, USA

E-mail address: rhn@math.umd.edu

(E. Otárola) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA, VALPARAÍSO, CHILE

E-mail address: enrique.otarola@usm.cl

(A.J. Salgado) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TENNESSEE, KNOXVILLE, TN 37996, USA

E-mail address: asalgad1@utk.edu